



ANTIBACTERIAL POTENTIAL OF CERTAIN AROMATIC OILS OF PLANT ORIGIN AND VACCINE DESIGN AGAINST METHICILLIN RESISTANT STAPHYLOCOCCUS AUREUS

SHUCHI KAUSHIK

Department of Biotechnology, College of Life Sciences,
Cancer Hospital & Research Institute, Gwalior (M. P.), INDIA.

Abstract:

Objective- "Staphylococcus" is a common inhabitant of the skin and mucous membrane and it accounts for a considerable proportion of human infections. Emergence of Multi Drug Resistance indicates a dire to understand the bacterial involvement in infections and find out new alternative approaches in its therapeutics and prevention. The antimicrobial properties of volatile aromatic oils derived from edible plants have been recognized since antiquity. An advantage of essential oils over antibiotics may be that they are classified as GRAS (Generally Recognized As Safe). The present study was undertaken to study the antimicrobial resistance patterns of *S. aureus* isolated from various samples collected from Hospitals of Gwalior. During the present study an effort was made to find out the information about *mecA* protein of Staphylococcus and their conserved regions were analyzed in order to assess their antigenic potential.

Materials & Methods- In the present study, a total of 872 samples were collected and processed for MRSA screening. Conventional methods were used for the isolation and identification of bacteria. Thereafter, antibacterial property of 20 various drugs as well as aromatic compounds of 18 herbal plants was performed against multiple resistant Staphylococcus aureus (MRSA) isolates according to the guidelines of National Committee for Clinical Laboratory Standards (NCCLS). In silico prediction of vaccine candidates in *mecA* through bioinformatics approach was also performed. In this regard, we have done multiple sequence alignment of *mecA* sequences fetched from database. Analysis was done for finding out the potential of these conserved regions to work as B cell epitope & T- cell epitope by the use of ProPred & EpiJen server. Structure prediction of this sequence was done by PSIPRED.

RESULTS & DISCUSSION-

The study revealed that drug resistance pattern of MRSA isolates is increasing. But the major concern is the development of resistance against Vancomycin which is thought to be the most effective drug against Staphylococcus. In comparison to antibiotics, essential oils showed very good activity against the test bacteria with few exceptions. The essential oils of Clove and Cinnamon were found to be more active against the test organism. The results support folkloric usage of the studied plants and suggest that most active extracts can be subjected to isolation of the therapeutic antimicrobials and undergo further pharmacological evaluation. The ancient practice of vaccine designing was being evaluated in the present study for the development of preventive measure against MRSA infections. Multiple sequence alignment revealed various conserved regions in *mecA* protein. We predicted one such type of multi-epitope peptide which was having very good potential to induce B cell response and a very good candidate for binding to MHC II molecule. This data can be very helpful for generating antigenic candidate by wet lab researchers.

Title:ANTIBACTERIAL POTENTIAL OF CERTAIN AROMATIC OILS OF PLANT ORIGIN AND VACCINE DESIGN AGAINST METHICILLIN RESISTANT STAPHYLOCOCCUS AUREUS.Source:Golden Research Thoughts [2231-5063] SHUCHI KAUSHIK
yr:2012 vol:2 iss:6

KEYWORDS:

Multidrug Resistance, GRAS, essential oils, in silico, multiepitope peptide.

INTRODUCTION

The genus "Staphylococcus" is common inhabitant of the skin and mucous membrane and it accounts for a considerable proportion of human infections. Humans are a natural reservoir for Staphylococcus aureus. Approximately 20–30% of the general population is "staph carriers" (Heyman, 2004). Staphylococcus aureus has been reported as a major cause of community and hospital acquired infection. It is still one of the four most common causes of nosocomial infections, often causing post-surgical wound infections. The emergence of antibiotic resistance in microorganisms and their spread is threatening the medical community. This is particularly true in case of Staphylococcus aureus.

The resistance development in Staphylococcus aureus dates back to 1940s. Multiple drug resistance of Staphylococcus aureus is due to the presence of mecA gene coding for penicillin binding protein (PBP2a) with a low affinity for β -lactam antibiotics. This gene is carried on Staphylococcal Cassette Chromosome (SCC) mec, a unique mobile genetic element that harbors the methicillin resistant gene (mecA) and other antibiotic resistant determinants. Thus, it mediates drug resistance in Staphylococci (Zhang et al., 2005).

The increasing gain of resistance to available antimicrobials and side effects associated with the drugs have attracted the attention of scientific community towards the search and development of new cost effective drugs of natural and synthetic origin (Fine et al., 2000). In the present scenario, interest of biologists to find antimicrobial properties in aromatic plants especially essential oils is continually increasing. Some studies demonstrated that plant derived essential oils may be an effective alternative to overcome microbial resistance to conventional drugs (Takarada et al., 2004; Didry et al., 1994). Essential oils are complex mixtures of volatile secondary metabolites that mainly consist of mono and sesquiterpenes including carbohydrates, alcohols, aldehydes, ketones and ethers. These are responsible for both the fragrant and biological effects of aromatic medicinal plants (Salzer, 1977; Angioni et al., 2003; Senatore et al., 2004). So, we also tried to analyze whether application of essential oils can be used to design a therapeutic measure to address this problem. In comparison to antibiotics, essential oils showed very good activity against the test bacteria with few exceptions. Varieties of aromatic medicinal plants are tested for treating infectious diseases (Kalemba & Kunicka, 2003). These plants have been mentioned in different phytotherapy manuals due to their availability, fewer side effects and reduced toxicity. An important characteristic of essential oils and their constituents is their hydrophobicity, which enables them to permeate in the lipids of bacterial cell membranes & mitochondria, thus disturbing the structures and rendering them more permeable (Sikkema et al., 1994; Sikkema et al., 1995).

No effective vaccine is generally available that stimulates active immunity against staphylococcal infections in humans. A vaccine based on fibronectin binding protein induces protective immunity against mastitis in cattle and might also be used as a vaccine in humans. However, vaccine therapies represent a new and innovative approach in broadening the available clinical tools against the global health problem of community and healthcare-associated S. aureus bacterial infections. Moreover it is cardinal to focus over mecA protein to design a multi-epitope antigenic target specific against multidrug resistant Staphylococcus that may provide protection from variety of infections caused by this deadly bacterium.

The objective of this research was to evaluate the potential of drugs and essential oils on standard microorganism strains as well as multi-drug resistant bacteria, which were isolated from hospitals. Moreover, we investigated the possible vaccine candidate for the test organism using bioinformatics tools.

MATERIALS & METHODS

Micro-organisms used in this study were isolated from patients suffering from various infections. The present study comprised 872 cases among which 500 samples were taken from patients, 266 cases were from healthy and 106 samples were taken from environment. All these cases were selected out of the peoples attending the local hospitals of Gwalior and the interior environment of the hospitals' wards. The candidates for this study were selected randomly.

The control group was selected from the persons who were not having any infection or any other visual symptom of diseases. The subjects for this study were between 5 to 75 years of age. Patients were examined by the doctors. Data were recorded, analyzed and subjected to statistical analysis.

CULTURE MEDIA AND ANTIBIOTICS

Brain Heart Infusion agar and Mueller-Hinton agar were used for growing the organisms. Antibiotic discs used in the study are given in table no. 1. Test bacteria were characterized by their biochemical characteristics as well as by standard microbiological techniques. The identified bacterial pathogens were tested against 20 different antibiotics using disc diffusion method of Bauer et al, 1966. MRSA was determined by the reference broth micro-dilution methods using established National Committee for Clinical Laboratory Standards break points.

Eighteen essential oils obtained from Department of pharmaceutical Science, Jiwaji University, Gwalior, M.P were used in this study. These oils were selected on the basis of their use in traditional and conventional medicine.

SCREENING OF ESSENTIAL OILS FOR THEIR ANTIMICROBIAL POTENTIAL

Each bacterial culture, to be tested was streaked onto Brain Heart Infusion (BHI) non-inhibitory agar medium to obtain isolated colonies. Each colony was put into BHI broth medium. After 6 hrs of growth, microorganisms at a concentration of 10^6 cells/mL, were layered on the surface of Mueller-Hinton agar plates. Subsequently, filter paper discs (6 mm dia.) saturated with essential oils (40 μ L) was placed on the surface of each inoculated plate. In order to evaluate the efficiency of the methodology, 40 μ L of each essential oil was instilled simultaneously in the wells of another plate. The plates were incubated at 37 °C. After 24 h. it was possible to observe the inhibition zones. The diameter of the zones of complete inhibition was measured in millimetres with the help of antibiotic zone scale (HiMedia Pvt. Ltd., Mumbai, India). Bacterial growth with halos equal to or greater than 10 mm was considered susceptible to essential oils. Controls were maintained along with each test.

VACCINE DESIGNING USING BIOINFORMATICS TOOLS

After the analysis of therapeutic potential of various natural products against MRSA, an effort was made to analyze preventive measure against it. The development of vaccine against several diseases is a very ancient practice. Recently this practice is also going to be evaluated for the development of preventive measure against MRSA infections. Molecular pathogenesis of MRSA infections involve several phases, each of which may offer targets for immunological intervention.

So we performed in silico prediction of vaccine candidates of *mecA* through bioinformatics approach. In this context, we have done multiple sequence alignment of *mecA* sequences using database. Results & Discussion

The threat to the human population is that reservoirs of drug-resistant bacteria is abound. The rotational use of vancomycin in the treatment of life-threatening infections due to MRSA is of last resort to preserve the efficacy of this antibiotic. The spectra of high-level vancomycin resistance transposed into MRSA are one of the doomsday scenarios in infectious diseases. There has also been the call to consider wider use of combinations of antibiotics in primary therapy, not because of their Synergistic approaches where two components work together to neutralize a single target, but combinations of distinct antibiotic classes that work on different targets concurrently.

In the present study, *S. aureus* was indicated by the yellow halo produced around the colonies on Mannitol Salt Agar. This is caused by the ability of *S. aureus* (and not *S. epidermidis* or most other osmotolerant bacteria) to ferment mannitol to acids which is detected by a change of pH indicator from red to yellow (Leboffe and Pierce 2002). Mannitol salt agar is often a selective medium for *Staphylococcus* due to the high concentration of sodium chloride the agar contains.

The observed colonies were circular and pale to dark yellow in color. The diameter of the colonies was about 1 to 1.5 millimeters. Gram staining of various colonies grown on different plates showed that the bacteria were Gram positive cocci, and arranged in irregular, grape like clusters.

Even though pharmaceutical companies have produced a number of new antibacterial drugs over the years, resistance to these drugs by *Staphylococcus aureus* has increased manifold and has now become a global concern. In general, *Staphylococcus aureus* have the genetic ability to transmit and acquire resistance to drugs used as therapeutic agents. Therefore the identification of new effective antimicrobial agents is of paramount importance. Medicinal plants have long been investigated as the potential sources among new agents,

Results on the antibiotic sensitivity of MRSA isolates are presented in table No.2. Overall study indicated that the antibiotics Cefuroxime, Ciprofloxacin, Piperacillin, Linezolid, Teicoplanin, Co-Trimoxazole, Ofloxacin, Tetracycline, Amikacin, Cefaperazone Sulbactam and Vancomycin should be

used judiciously against most of the staphylococcal infections in the region. Ampicillin, Cefazolin, Chloramphenicol, Erythromycin, Gentamycin, Oxacillin and Penicillin, however, should not be used any more by general medical practitioner while giving treatment to the patients because organisms are resistant to these antibiotics.

This study also showed that antimicrobial resistance of *Staphylococcus aureus* was high and alarming. The major concern is the development of resistance against Vancomycin which is thought to be the most effective drug against *Staphylococcus*. Resistant bacteria could be the mutant form of common bacteria due to non-judicial use of broad spectrum antibiotics. Antibiotics are frequently prescribed in hospital and general practice. However they are often administered before the pathogen's culture and sensitivity results are known. Oxacillin resistance is primarily mediated by the production of penicillin binding protein 2a (PBP2a), which is encoded by the *mecA* gene. Because its production is under complex control by regulatory genes and in some cases also by *mecA*-negative resistance mechanisms, phenotypic or even genotypic evaluation is not completely straightforward. Furthermore, genetic testing requires significant technical and financial resources which might be lacking in many clinical laboratories. It is possible that plasmid DNA represents the transposon-associated *vanA* gene cluster found in vancomycin-resistant enterococci, which could be horizontally transferred to other species, consequently conferring resistance to the antibiotic in that species (Archer and Crossley 1997). It is also likely that strain's resistance to vancomycin was a result of a thickened cell wall, which allows the bacteria to inhibit vancomycin molecules from reaching its major target on the cytoplasmic membrane of the cell (Cui and Hiramatsu 2003). Plasmid borne antibiotic resistance was beyond the scope of the present study.

As the distribution of causative organisms and bacterial resistance rates vary according to time and place, the recent local data will be conducive to the clinicians for the best choicest treatment. As a result, not only the patients will be treated with the correct antibiotics but the misuse and overuse of antibiotics, which lead to rapid development and spread of resistance, will be minimized.

Since the development of antibiotic resistance in the pathogenic strains of *Staphylococcus aureus* is an ever increasing problem, suitable and possible alternate chemotherapeutic phytochemical compounds of plant origin such as alkaloids, terpenoids, polyphenols, flavonoids and steroids may be tried for effective control of drug resistant bacteria like MRSA (Prakash et al., 2006).

The antibacterial activity of essential oils of 15 plant species was assayed in vitro by disc diffusion and agar well diffusion method against the test bacteria. The maximum antibacterial activity was shown by *Cinnamomum zeylanicum*, followed by clove. The essential oils of olive, mustard, castor and coconut could not inhibit any of the bacterial isolates studied. It is quite possible that some of the essential oils that are ineffective in this study do not possess antibacterial properties or the essential oils might have contained antibacterial constituents in insufficient concentration so as to be effective. The activity of cinnamon oil was found to be more than vancomycin drug. So this can be a better agent for inhibition of MRSA after suitable toxicological analysis. All other oils showed moderate activity.

Cinnamon oil and clove oil are very ancient ingredients of Indian food. Their antimicrobial activity further supports the traditional natural medicinal system of India. Previous GC-MS study of cinnamon oil has shown Cinnamyldehyde as the predominant active component (Simic et al., 2004; Barrata et al., 1998). Cinnamyldehyde is a natural anti-oxidant and the animal studies suggest that an extract of cinnamon bark taken orally may help to prevent stomach ulcer. Cinnamyldehyde was completely inhibiting both sensitive and resistant strain of *Helicobacter pylori* (Ali et al., 2005). Cinnamon oil was not harmful when consumed in food products and it inhibited the growth of molds, yeast and bacteria (Matan et al., 2006). Cinnamon extract had a regulatory role in blood glucose level and lipids. This oil is also used in the treatment of cancer and other microbial diseases (Nadkarni, 1976). It can be incorporated into creams, lotions, drops, etc. which are applied externally on the body to treat diseases caused by *Aspergillus niger*.

Clove oil has also been shown as potent antimicrobial agent against various pathogenic strains of bacteria (Islam et al., 1990) and fungi including *Aspergillus*, *Penicillium* and yeast etc. (Matan et al., 2006). The main constituents of the clove oil are phenyl propanoids such as carvacrol, thymol, eugenol and cinnamaldehyde (Chaieb et al., 2007). The characteristic aroma of clove is due to the compound eugenol. It is the main component of clove essential oil, comprising 72-90% (Bensky et al., 2004). This can be helpful in designing antimicrobial agents for topical and oral applications.

The use of plants to heal diseases, including infectious one, has been extensively employed by people. Data from the literature as well as our own findings reveal that the plants have the great therapeutic potential, in spite of the fact that they have not been completely investigated. Therefore, more studies need to be conducted to search newer compounds. Once extracted, and before being used in new treatments, they should have their in vivo toxicity tested.

Furthermore, in a few cases, these plant aromatic oils were found active against antibiotic resistant bacteria in a very low concentration with minimum possible toxic effects, and can be used in the treatment

of infectious diseases caused by the resistant microbes.

In the present study, it was found that *mecA* protein of *Staphylococcus* can also be used as an effective candidate for the development of preventive measures against drastic diseases by blocking its resistance efficiency. In fact in silico approach the vaccine target prediction is definitely less labour intensive, rapid and economic in relation to search for a lead antigenic molecule against *mecA* protein. Multiple sequence alignment revealed various conserved regions of this protein. The region found to be conserved in all *mecA* of *Staphylococcus* studied was

Analysis was done for finding out the potential of these conserved regions to work as B cell epitope and we found 41 potential B cell epitopes among the conserved sequence. For any predicted epitope, it is cardinal that it should induce T and B cell response. Such type of multi-epitope vaccine is a very recent experimental technique for predicting vaccine targets against HIV and Influenza virus (). We predicted one such type of multi-epitope peptide which was having very good potential to induce B cell response as well as a very good candidate for binding to MHC II molecule. Among 51 MHC II alleles analyzed, the conserved region of protein *mecA* showed binding affinity with all the 51 alleles with one or more than one paratope found in each. As per the results of ProPred, 47 sequences were found to bind with MHC I alleles. During the last step of study, this protein sequence was also analyzed for its binding affinity with T-cell receptors using the EpiJen server. Structure prediction of this sequence by PSIPRED revealed that 22 helix are present in this sequence. This data can be very helpful for generating antigenic candidate by wet lab researchers.

The results presented in this report were encouraging, although clinical controlled studies are required to define the real efficacy and possible toxic effects in vivo. It is also visualized that there is a need for extensive investigation concerning the molecular basis of understanding the mechanism which is fundamental in the development of pharmacological agents of medicinal plants used in the treatment of bacterial infections.

ACKNOWLEDGEMENTS

I am grateful to Dr Archana Shrivastav, Dr Sudhir Kumar Jain and Dr Asha Mukul Jana for their comments and suggestions. This study was conducted with the kind permission of Sh. Shitla Sahai ji, founder Trustee of Cancer Hospital and Research Institute, Gwalior, M.P. I am also thankful to Dr. B R Shrivastav, Director of Cancer Hospital and Research Institute, Gwalior for his kind support.

REFERENCES

1. Heyman, D. Control of Communicable Diseases Manual (2004) 18th Edition. Washington DC: American Public Health Association.
2. Zhang, K., McClure, J., Elsayed, S., Louie, T., Conly, J. M. 2005. Novel multiplex PCR Assay for characterization and concomitant subtyping of staphylococcal cassette chromosome *mec* types I to V in methicillin resistant *Staphylococcus aureus*. *J Clin Microbiol.* 43, 5026-5033.
3. Fine DH, Furgang D, Barnett ML, Drew C, Steinberg L, Charles CH, Vincent JW (2000). Effect of an essential oil-containing antiseptic mouthrinse on plaque and salivary *Streptococcus mutans* levels. *J Clin Periodontol* 27(3):157-61.
4. Takarada K, Kimizuka R, Takahashi N, Honma K, Okuda K, Kato T (2004) A comparison of the antibacterial efficacies of essential oils against oral pathogens. *Oral Microbiol Immunol* 19: 61-64.
5. Didry N, Dubreuil L, Pinkas M (1994). Activity of thymol, carvacrol, cinnamaldehyde and eugenol on oral bacteria. *Pharm Acta Helv* 69:25-28.
6. Salzer UJ (1977). The analysis of essential oils and extracts (oleoresins) from seasonings-a critical review. *CRC Crit Rev Food Sci Nutr* 9: 345-373.
7. Angioni A, Barra A, Arlorio M, Coisson JD, Russo MT, Pirisi FM, Satta M, Cabras P (2003). Chemical composition, plant genetic differences, and antifungal activity of the essential oil of *Helichrysum italicum* G. Don ssp. *microphyllum* (Willd) Nym. *J. Agric Food Chem.* 51: 1030-1034.
8. Senatore F, Arnold NA, Piozzi F (2004). Chemical composition of the essential oil of *Salvia multicaulis* Vahl. var. *simplicifolia* Boiss. growing wild in Lebanon. *J. Chromatogr. A.* 1052: 237-240.
9. Bauer, A.W., Kirby, W.M.M., Sherris, J.C. & Turck, M. (1966). Antibiotic susceptibility testing by a standardized single disk method. *American Journal of Clinical Pathology* 45, 493-6.
10. Leboffe, M.J. and B.E. Pierce. (2002). *Microbiology: Laboratory Theory and Application*. Englewood, CO: Morton Publishing Company.
11. Archer, G.L. and K.B. Crossley (Eds.). (1997). *The Staphylococci in Human Disease*. New York: Churchill Livingstone Inc.

12. Cui, L. and K. Hiramatsu. (2003). Vancomycin-Resistant *Staphylococcus Aureus*. In A.C. Fluit & F.J. Schmitz (Eds.), *MRSA Current Perspectives* (pp. 187-212). Norfolk, England: Caister Academic Press.
13. Prakash, M., V. Karthikeyan, S. Karuppusamy and N. Karmegam, 2006. Synergistic activity of certain plant extracts against Methicillin resistant *Staphylococcus aureus* (MRSA). *Journal of Ecotoxicology and Environmental Monitoring*, 16: 387-389
14. Kalembe D and Kunicka A (2003). Antibacterial and antifungal properties of essential oils. *Curr Med Chem* 10: 813-829.
15. Sikkema J, de Bont JA, Poolman B (1994). Interactions of cyclic hydrocarbons with biological membranes. *J Biol Chem* 269: 8022-8028.
16. Sikkema J, de Bont JA, Poolman B (1995). Mechanisms of membrane toxicity of hydrocarbons. *Microbiol Rev* 59: 201-222.
17. Matan N, Rimkeeree H, Mawson AJ, Chompreeda P, Haruthaithanasan V, Parker M (2006) Antimicrobial activity of cinnamon and clove oils under modified atmosphere conditions. *Int J Food Microbiol* 2006, 107:180-185.
18. Aureli P, Costantini A, Zolea S (1992) Antibacterial activity of some plant essential oils against *Listeria monocytogenes*. *J Food Prot* 55:344-348
19. Chopra RN, Nayar SL, Chopra IC (1956). *Glossary of Indian medicinal plants*. Council of Scientific and Industrial Research, New Delhi.
20. Prasad S, Mitra R (1970). Pharmacognostical studies on Ransa (*Pluchea lanceolata* C.B. Clarke and Olive). *J Res Ind Med* 5 (1): 59
21. Simic A, Sokovic MD, Ristic M, Grujic-Jovanovic S, Vukojevic J, Marin PD (2004) The chemical composition of some Lauraceae essential oils and their antifungal activities. *Phytother Res*, 18:713-717.
22. Baratta MT, Dorman HJ, Deans SG, Figueiredo AC, Barroso JG, Ruberto G (1998) Antimicrobial and antioxidant properties of some commercial essential oils. *Flav Fragr J*, 13:235-244.
23. Ali SM, Khan AA, Ahmed I, Musaddiq M, Ahmed KS, Polasa H, Venkateswar Rao L, Habibullah CM, Sechi LA, Ahmed N (2005) Antimicrobial activities of Eugenol and Cinnamaldehyde against the human gastric pathogen *Helicobacter pylori*. *Annals of Clinical Microbiology and Antimicrobials*, 4:20.
24. Kim SH, Hyun SH, Choung SY (2006) Anti-diabetic effect of cinnamon extract on blood glucose in db/db mice. *J Ethnopharmacol*, 104:119-123.
25. Quale JM, Landman D, Zaman MM, Burney S, Sathe SS (1996) In vitro activity of *Cinnamomum zeylanicum* against azole resistant and sensitive *Candida* species and a pilot study of cinnamon for oral candidiasis. *Am J Clin Med*, 24:103-109
26. Nadkarni KM (1976) *Indian Materia Medica*, Popular Prakashan, Bombay, India, pp.228-231.
27. Chaieb K, Hajlaoui H, Zmantar T, Ben Kahla-Nakbi A, Rouabhia M, Mahdouani K, Bakhrouf A (2007) The chemical composition and biological activity of clove essential oil, *Eugenia caryophyllata* (*Syzygium aromaticum* L. Myrtaceae): a short review *Phytotherapy Research*, 21(6): 501 – 506
28. Bensky D, Clavey S, Stoger E, Gamble A, Lai Bensky L (2004) *Chinese Herbal Medicine: Materia Medica*, (III Ed.) Eastland Press. *Beurteilung. Dtsch Zahnarzt Z* 3:189-191.