

Vol 3 Issue 11 May 2014

ISSN No :2231-5063

International Multidisciplinary Research Journal

Golden Research Thoughts

Chief Editor
Dr.Tukaram Narayan Shinde

Publisher
Mrs.Laxmi Ashok Yakkaldevi

Associate Editor
Dr.Rajani Dalvi

Honorary
Mr.Ashok Yakkaldevi

Welcome to GRT

RNI MAHMUL/2011/38595

ISSN No.2231-5063

Golden Research Thoughts Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

International Advisory Board

Flávio de São Pedro Filho Federal University of Rondonia, Brazil	Mohammad Hailat Dept. of Mathematical Sciences, University of South Carolina Aiken	Hasan Baktir English Language and Literature Department, Kayseri
Kamani Perera Regional Center For Strategic Studies, Sri Lanka	Abdullah Sabbagh Engineering Studies, Sydney	Ghayoor Abbas Chotana Dept of Chemistry, Lahore University of Management Sciences[PK]
Janaki Sinnasamy Librarian, University of Malaya	Catalina Neculai University of Coventry, UK	Anna Maria Constantinovici AL. I. Cuza University, Romania
Romona Mihaila Spiru Haret University, Romania	Ecaterina Patrascu Spiru Haret University, Bucharest	Horia Patrascu Spiru Haret University, Bucharest,Romania
Delia Serbescu Spiru Haret University, Bucharest, Romania	Loredana Bosca Spiru Haret University, Romania	Ilie Pinteau, Spiru Haret University, Romania
Anurag Misra DBS College, Kanpur	Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Xiaohua Yang PhD, USA
Titus PopPhD, Partium Christian University, Oradea,Romania	George - Calin SERITAN Faculty of Philosophy and Socio-Political Sciences AL. I. Cuza University, IasiMore

Editorial Board

Pratap Vyamktrao Naikwade ASP College Devrukh,Ratnagiri,MS India	Iresh Swami Ex - VC. Solapur University, Solapur	Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur
R. R. Patil Head Geology Department Solapur University,Solapur	N.S. Dhaygude Ex. Prin. Dayanand College, Solapur	R. R. Yaliker Director Managment Institute, Solapur
Rama Bhosale Prin. and Jt. Director Higher Education, Panvel	Narendra Kadu Jt. Director Higher Education, Pune	Umesh Rajderkar Head Humanities & Social Science YCMOU,Nashik
Salve R. N. Department of Sociology, Shivaji University,Kolhapur	K. M. Bhandarkar Praful Patel College of Education, Gondia	S. R. Pandya Head Education Dept. Mumbai University, Mumbai
Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai	Sonal Singh Vikram University, Ujjain	Alka Darshan Shrivastava Shaskiya Snatkottar Mahavidyalaya, Dhar
Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College, Indapur, Pune	G. P. Patankar S. D. M. Degree College, Honavar, Karnataka	Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore
Awadhesh Kumar Shirotriya Secretary,Play India Play,Meerut(U.P.)	Maj. S. Bakhtiar Choudhary Director,Hyderabad AP India.	S.KANNAN Annamalai University,TN
	S.Parvathi Devi Ph.D.-University of Allahabad	Satish Kumar Kalhotra Maulana Azad National Urdu University
	Sonal Singh, Vikram University, Ujjain	

Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.aygrt.isrj.net



ISOLATION OF HEAT RESISTANT FUNGI FROM CANNED FRUITS

Sani Ibrahim , M.B.Anusha* and P. Udhayaraja**

*Assistant Professor, Department of Biotechnology, Faculty of Science and Humanities, SRM University, Kattankulathur.
**Assistant Professor, Department of Microbiology, Selvamm Arts and science college, Namakkal.

Abstract:-Increase in the consumption of canned fruit juice rather than the fruit itself raises alarm to the safety of these canned juices as they are stored for longer period of time unlike the fruit and dormant spores present could germinate and cause spoilage. Heat-resistant molds are characterized by the production of ascospores or similar structures with heat resistance, in some instances comparable to bacterial spores. This enables them to survive even when they are thermally processed. So the presence of heat resistant molds in canned fruits is worthy of studying. In this research, heat resistant moulds exist in canned fruit juice and they can withstand the heat treatment/ pasteurization done at processing plants. Their presence might be due to contamination right from the farm fields or the processing company. Heat resistant moulds in canned fruit juice can pose a set or health hazard in the body which includes Diarrhea and Stomachache.

Keywords: Heat resistant molds, Contamination, antibiotic etc.

INTRODUCTION :-

Heat resistant molds can survive heat treatments used for fruit processing and can grow and cause spoilage during storage. The heat resistance is attributed to the formation of sexual spores called ascospores and the heat resistance of the ascospores depends on the species, strain, pH, heating medium and other growth conditions (Beuchat and Pitt, 2006). Spoilage of fruits and fruit products by heat resistant molds has been recognized in several countries (Beuchat, 1978). *Byssosclamyces fulva*, *B. nivea*, *Neosartorya fischeri*, *Talaromyces macrosporus*, *T. bacillisporus*, and *Eupenicillium feldianum* have been most frequently encountered. These organisms are saprophytic rather than parasitic and usually contaminate fruits on or near the ground. *Byssosclamyces* species have been recognized as spoilage molds in fruit since the early 1930s and have been extensively studied (Hull, 1979). Spoilage by other heat-resistant molds is a less serious problem recognized only in recent years. Consequently, less information is available concerning the behavior of these other genera in thermally processed fruit products (Beuchat and Pitt, 2006). Germination of ascospores may result in visible growth of mycelia on fruits and fruit products. Production of pectic enzymes by *Byssosclamyces* can result in complete breakdown of texture in fruits and also can result in off-flavor development. Heat-resistant molds, therefore, may constitute a public health hazard as well as spoilage problems (Put and Kruiswijk, 1984). Some *Byssosclamyces* species produce patulin, byssotoxin A, and byssochlamic acid, all having toxic effects on laboratory animals. *Neosartorya fischeri* is known to produce fumitremorgin A, B, and C, terrein, and verruculogen (Nelson, et al., 1988).

OBJECTIVES

The aim of this research is to assess the presence or otherwise of heat resistant molds in canned fruit juice.

1. To isolate and identify the heat resistant molds in selected fruit juices.
2. To find the opinion of consumers regarding the contamination of the canned fruit juices.

HYPOTHESIS:

(i) Heat resistant molds are present in canned fruit juice.

MATERIALS AND METHOD

The method described by Murdock and Hatcher (1978) and modified by Beuchat and Pitt (2006) was used in this research.

SAMPLE COLLECTION

The samples of this research are canned Orange, Mango, Pineapple, Apple and Tomato juice. The Samples being already in liquid form hence no blending was required.

MYCOLOGICAL ANALYSIS

Malt extract agar (MEA) was used as the culture media, and it was prepared according to the manufacturing guidelines/instructions. Growth of bacterial spores that may survive the heat shock has been prevented by the inclusion of an antibiotic (erythromycin) as described by Pitt and Hocking, (1997). Finally it was sterilized by autoclaving at 121°C for 15 minutes.

Pour plate method was used in preparation of plates for incubation. Five (5) plates were prepared for each fruit; 2ml of each sample was carried by using a sterilized 5ml syringe and transferred into a petridish. Malt extract media (prepared) was then poured into each plate; the content is then mixed gently by shaking the plates clockwise then counter-clockwise. It was allowed to solidify on a labeled surface to obtained even distribution, the plates then were inverted.

The plates were incubated at 30°C in an incubator machine for seven (7) days, the control samples were like-wise incubated. After the incubation period (7 days), each colony was then morphologically studied and recorded.

IDENTIFICATION

The identification of heat resistant molds was done according to the procedure described by Beuchat and Pitt (2006). And it was achieved by morphological examination of colonies formed, and by microscopic examination. The size of the colonies were measured by using ruler and recorded. Hand magnifier was used to aid view. The Isolates were Identified by making reference to the taxonomic key described by Beuchat and Pitt (2006).

MICROSCOPIC EXAMINATION

A wet mounts was prepared by putting a small piece of fungal filament using sterilized inoculating needle in a clean free-grease glass slide. A drop of lactophenol was added for clear view. This was then covered by cover slip and examine under x40 by x100 objective lens. The mycelium was observed i. e whether the hyphae are coenocytic or septate, branched or unbranched. The arrangement of spores were observed and also recorded. Also the diversity was recorded by grouping the species into different taxonomic genera.

RESULTS

A total of 31 fungal isolates were obtained from the five canned fruit samples screened. The isolate were classified by grouping into three taxonomic genera namely; *Byssochlamys*, *Talaromyces* and *Neosartorya*. Table A shows the average diameter covered by the isolates on the plates with the highest mean recorded in Orange and least in Apple. Table B showed the isolates and their respective frequency on the plates, the highest number of isolates was also recorded in Orange juice. Table C shows the percentage of respondents that encounter changes in their fruit juices and those that still consume the fruit juices when changes are noticed. Table D shows the percentages of the respondent's perception on the cause and source of contamination. Table E shows the percentages of the respondent's fruit juice which contamination occur most and the effect on health when consumed. All results are presented below:

Table A
SHOWING AVERAGE DIAMETERS COVERED BY THE ISOLATES ON THE PLATES

PLATE NUMBER	TOMATO	ORANGE	MANGO	PINEAPPLE	APLLE
1	2.1	2.5	3.0	2.9	0.0
2	2.4	3.3	2.8	4.1	0.0
3	2.0	4.6	4.4	3.6	2.0
4	3.0	2.4	1.9	4.3	2.2
5	3.0	2.9	0.0	0.0	2.5
TOTAL	12.5	17.7	12.1	14.9	6.7
MEAN	2.5	3.5	2.4	3.0	1.3

TABLE B
SHOWING THE ISOLATES AND THEIR FREQUENCY ON THE PLATES

Fruits	Isolates	Frequency	Percentage
Orange	<i>Byssochlamysnivea</i>	3	33.3%
	<i>Byssochlamysfulva</i>	3	33.3%
	<i>Neosartoryafisherii</i>	3	33.3%
Control	-	-	-
Mango	<i>Neosartoryafisherii</i>	2	25%
	<i>Byssochlamysfulva</i>	5	62.5%
	<i>Byssochlamysnivea</i>	1	12.5%
Control	-	-	-
Apple	<i>Talaromycesmacrosporuos</i>	3	100%
Control	-	-	-
Tomato	<i>Talaromycesmacrosporuos</i>	2	33.3%
	<i>Neosartoryafisherii</i>	4	66.7%
Control	-	-	-
Pineapple	<i>Byssochlamysfulva</i>	2	40%
	<i>Byssochlamysnivea</i>	3	60%
Control	-	-	-

TABLE C
SHOWING THE PERCENTAGES OF RESPONDENTS THAT ENCOUNTER CHANGES IN FRUIT JUICE AND THOSE THAT STILL CONSUME THE JUICE AFTER THE CHANGES ARE NOTICED.

CHANGES ENCOUNTERED	PERCENTAGE	CONSUMPTION AFTER POILAGE	PERCENTAGE
Yes	66.25%	Yes	19.72%
No	33.75%	No	80.28%

TABLE D
SHOWING THE PERCENTAGES OF THE RESPONDENTS PERCEPTION ON THE CAUSE AND SOURCE OF CONTAMINATION RESPECTIVELY.

CAUSE OF SPOILAGE	PERCENTAGE	SOURCE OF CONTAMINATION	PERCENTAGE
Virus	0%	Farm fields	30.26%
Bacteria	35.21%	Processing company	43.42%
Fungi	64.79%	Retail shops	26.37%

TABLE E
SHOWING THE PERCENTAGES OF THE RESPONDENT'S FRUIT JUICE WHERE CONTAMINATION OCCUR MOST AND THE EFFECT ON HEALTH WHEN CONSUMED

MOST ENCOUNTERED FRUIT JUICE	PERCENTAGE	EFFECT ON HEALTH	PERCENTAGE
Orange	25%	Stomachache	36.21%
Mango	25%	Diarrhea	15.52%
Pineapple	12.5%	Both stomachache and diarrhea	12.07%
Tomato	35.94%	None	29.31%
Apple	1.56%	Other effects	6.90%

DISCUSSION

The result of this experiment showed that, three genera of heat resistant molds were isolated and identified namely; *Byssochlamys*, *Talaromyces*, *Neosartorya*, out of the five plates of orange cultured. There was growth on two of the tested samples. This showed the occurrence or contamination of heat resistant molds in orange and tomato samples, translating to a total of 17.7cm and 12.5cm total diameter and 3.5cm and 2.5cm mean diameter for orange and tomato juice respectively. Apple juice sample has the least with only three positive plates and a total diameter of 6.7cm and 1.3cm mean diameter covered on the plates.

Stolk and Samson (1981) observed the presence of heat resistant mould in canned fruit juice samples, including all the five used in this study and confirmed the presence of different species including *Byssochlamys* sp, *Talaromyces* sp, *Neosartorya* sp. Contamination of fruit during harvest as well as growth of fungi on lesions and wounds have been described by Moyer et al., (1969) as a major cause of contamination in canned fruit juice. Presence of heat resistant moulds in pipes of processing companies is also a source of contamination according to Murdock and Brokaw (1958).

Byssochlamys was described by Beuchatt and Pitt (2006) as the most economic important heat resistant mould in canned fruits and this may be due to the production of byssochlamic acid which breaks down the fruit structure. They also have

toxic effect. According to Beuchat and Pitt (2006), *Byssochlamysfulva* produce colonies that cover the whole Petridish and relatively sparse, with low or somewhat floccose, with conidia production heavy, uniform and colored olive brown, while *B. nivea* produce white to slightly gray colonies on malt extract agar. Therefore the colonies formed can be confirmed to be *B. fulva* and *B. nivea*. This is in agreement with the results of this research.

Byssochlamys spp. was first reorganized as a spoilage agent in Great Britain by Oliver and Rendle (1934) and for a time it appeared to be a localized problem. However in recent years, spoilage outbreaks due to this mould has been recognized in almost all the continents of the world (Maunder, 1989) The occurrence of *Neosartorya* spp on orange sample only was due to the facts that, this species grow only on food with low pH. Orange has a pH of 1.8-2.9 while that of tomato, pineapple and apple have a range of 2.5-3.7, 3.2-4.4 and 3.2-4.5 respectively, *Talaromyces* spp usually grows in pH range of 2.9-3.3 and therefore are more likely to be present in high rate in tomato and apple than the other two samples (Maunder, 1989).

Due to high concentration of sugar (carbohydrate) in orange, *Byssochlamys* spp are found to be more abundant. *Byssochlamys* spp grow best at high concentration of sugars (Norman, 1980). The occurrence of these organism may be attributed to their ability to produce resistant spores, *Byssochlamys* spp more resistant to high temperature. In conjunction to this, *Byssochlamys* spp has been implicated in the spoilage of fruits and vegetables in Nigeria (Uzuegbu and Emiforiye, 1984). All the samples showed contamination by these heat resistant molds. This may be due to improper handling right from field or during transportation or in during processing. (Pitt and Hocking, 1985).

RECOMMENDATIONS

1. It is also recommended by virtue of this research that canned fruit juices should be stored at low temperatures as high temperature could activate the ascospores of the molds and cause spoilage.
2. Contaminated juices should not be consumed as they may pose health hazard.
3. Processing companies should ensure that all their fruits have bearable or minimum concentration of contaminants when relieved from the farm and they should employ better processing techniques such as flash pasteurization which ensures better safety.

CONCLUSION

It can be concluded from the results of this research that heat resistant moulds exist in canned fruit juice and they can withstand the heat treatment/ pasteurization done at processing plants. Their presence might be due to contamination right from the farm fields or the processing company. Heat resistant moulds in canned fruit juice can pose a set or health hazard in the body which includes Diarrhea and Stomachache.

REFERENCES

1. Balla C Farkas J. (2006). Minimally processed fruits and fruit products and their microbiological safety. In: Hui YH, Barta J, Cano MP, Gusek T, Sidhu JS, Sinha N, editors. Handbook of fruits and fruit processing. Anies, Iowa: Blackwell Publishing, p 115-28.
2. Bayne, H. G. and Michener H. D., (1976). Heat resistance of *Byssochlamys* ascospores. Applied environmental microbiology. 37: 449 - 453.
3. Beuchat, L.R, Pitt J.L (1992). Detection and enumeration of heat-resistant molds. In: Vanderzant C, Splittstoesser DF, editors. Compendium of methods for the microbiological examination of food. Washington, D.C.: American Public Health Assoc. p 251-63.
4. Beuchat, L.R. (2002). Ecological factors influencing survival and growth of human pathogens on raw fruits and vegetables. Microb Infect 4:413-23.
5. Beuchat, L. R. And Pitt J.I. (2006). Detection and enumeration of heat resistant molds. In compendium of methods for the microbiological examination of foods. 2nd edition, American public health association, Washington d. C. 21: 217 - 222.
6. Beuchat, R.L (1978). Beuchat Food and Beverage Mycology. AVI Publishing Comp. INC. Westport, Connecticut.
7. Brackett, R.E. (1994). Microbiological spoilage and pathogens in minimally processed refrigerated fruits and vegetables. In: Wiley RC, editor. Minimally processed refrigerated fruits and vegetables. New York: Chapman & Hall, p 269-312.
8. Brackett, R.E. (2001). Fruits, vegetables and grains. In: Doyle MP, Beuchat LR, Montville TJ, editors. Food microbiology. Fundamentals and frontiers. 2nd edition. Washington, D.C.: ASM Press, p 127-38.
9. Chang, S.S, Kang, DH. (2004). *Alicyclobacillus* spp. in the fruit juice industry: history, characteristics, and current isolation/detection procedures. Crit Rev Microbiol 30:55-74.
10. Coursey, D.G., Burden A. and Richard J. E., (1976) Recent Advances in Research on post harvest Handling of Tropical and subtropical fruits. Acta Horticulture 57:135-143.
11. Davenport, C. F and Beech, H. (1970). Role of yeast in fruits spoilage. Phytopathal review. Vol. 13, pp22.
12. Harris L.J, Farber J.N, Beuchat L.R, Parish M.E, Suslow T.V, Garrett E.H, Busta F.F. (2003). Outbreaks associated with fresh produce: incidence, growth, and survival of pathogens in fresh and fresh-cut produce. CRFSFS 2(1):78-141.
13. Hull, R. (1979). Study of *Byssochlamysfulva* And Control Measure in Processed fruit. Journal of applied biology 26:800-

822.

14. Igbeka J.O. and Ike A. (1982). Stimulation of moisture movement during drying. *Journal of food technology*. 17:27-36.
15. James, M.J., (1978). Natural antimicrobials in postharvest storage of fresh fruits and vegetables.
16. In: Roller S, editor. Natural antimicrobials for the minimal processing of foods. Boca Raton, Modern food microbiology. 3:38-55. Jay J.M, Loessner M. J, Golden D.A. (2005). Modern food microbiology. 7th edition, New York:Springer Science Business Media Inc. 790 p.
17. John, F.D. (1979). Studies of mucor causing post harvest decay of fresh fruits product. 69:865-869
18. Jones, A. L. and Aldwinckle, H.S (1990). Compendium of apple and pear diseases. American phytopathological society press, st. Paul, mn.
19. Kalia A, Gupta R.P. (2006). Fruit microbiology. In: Hui YH, editor. Handbook of fruits and fruit processing. Ames, Iowa: Blackwell Publishing, p 3-28.
20. King, A.D. (1969). Related heat resistant fungi in grape products. *Applied microbiology*. 18: 166-173
21. Kingsley R. S. (2008), Microsoft Encarta, 2008. Microsoft Corporation, USA.
22. Lund, B.M, Snowdon, A.L. (2000). Fresh and processed food. In: Lund BM, Baird-Parker TC, Gould GW, editors. The microbiological safety and quality of food. Gaithersburg, Md.: Aspen Publication, p 738-58.
23. Mart'nez A, D'laz R.V, Tapia, M.S. (2000). Microbial ecology of spoilage and pathogenic flora associated with fruits and vegetables. In: Alzadora SM, Tapia MS, L'opez-Malo A, editors.
24. Minimally processed fruits and vegetables. Fundamental aspects and applications.
25. Gaithersburg, Md.: Aspen Publishers, Inc. p 43-62. Martinez, J. (1949). Temperature tolerance of *Aspergillus fischeri* var. *glaber* in canned strawberries. *Irish J. Agric. Res.* 9:59-67.
26. Maunder, D: T. (1989). Spoilage problems caused by moulds *Byssoschlamys*- *Paecilomyces* group seminar Abstracts. *Circ.* 20. N.T. State Agric. Exp. Stn., Geneva.
27. Montville TJ, Matthews KR. (2001). Principles which influence microbial growth, survival, and death in food. In: Doyle MP, Beuchat LR, Montville TJ, editors. Food microbiology: fundamentals and frontiers, 2nd edition. Washington, D.C.: American Society for Microbiology Press, p 13-32.
28. Moreau, C. (1979). Pathogen reduction in unpasteurized apple cider: adding cranberry juice to enhance the lethality of warm hold and freeze-thaw steps. *J Food Prot* 69:293-8.
29. Mukhtar, F. B. (2008). An Introduction to Biostatistics. Spectrum books limited, Ibadan. pp 61-62
30. Murdock D. I and Brokaw C. H (1958) Sanitary control in processing citrus concentration. *Food technol.* 12, 573-576
31. Murdock, D. L., and W. S. Hatcher, Jr. 1978. A simple method to screen fruit juices and concentrates for heat-resistant mold. *J. Food Prot.* 41:254-256.
32. Naidu A.S. (2000). Lactoperoxidase. In: Naidu AS, editor. Natural food antimicrobial systems. Boca Raton, Fla.: CRC Press, p 103-32.
33. Nguyen-The C, Carlin F. (1994). The microbiology of minimally processed fresh fruits and vegetables. *Crit Rev Food Sci Nutr* 34:371-401.
34. Nelson, P. V., L. R. Beuchat, and J. C. Frisvad. (1988). Growth and fumitremorgin production by *Neosartorya fischeri* affected by temperature, light, and water activity. *Appl. Environ. Microbiol.* 54:1504-1510.
35. Norman, W.D (1980). Influence of relative humidity conditions on the thermal resistance of several kinds of spores of molds. *Acta Aliment.* 2:163-171.
36. Oliver, M I. and Rendle, T. (1934). A new problem in fruit preservation Studies on *Byssoschlamys fulva* and its effect on tissues of processed fruit. *J. soc-chem. Ind. London* 53, 166-172.
37. Oliver, M. (1934). A new problem in fruits and its effect on the tissue of processed fruits. *Journal of social chemical industries.* 53:166t-172t.
38. Pantastico, B. (1975). Post harvest physiology, handling and utilization of tropical and sub tropical fruits and vegetables. Pp 415, 424 and 426.
39. Pao S, Petracek P.D. (1997). Shelf-life extension of peeled oranges by citric acid treatment. *Food Microbiol* 14:485-91.
40. Pitt, J. I and Hocking, (1997). Fungi and food spoilage, 2nd edition. Blackie academic and professional, London.
41. Put, H.M. and Kruiswijk J.T., (1984). Disintegration and organoleptic deterioration of processed strawberries caused by the molds *B. nivea*. *Journal of applied bacteriology.* 27:53-58
42. Reid, W. W (1992). The pectic enzymes of the fungus *B. fulva*. *Journal of biochemistry* 4:57-77 Roger, Y. S (1983). Incidence of *B. fulva* in queen land grown in canned fruits. 22:247-350
43. Roland, J. O, Beuchat, L. R. (1984). Biomass and patulin production by *Byssoschlamys nivea* in apple juice as affected by sorbate, benzoate, SO₂ and temperature. *J Food Sci* 49:402-6.
44. Ross A.I., Griffiths, M.W, Mittal, G.S, Deeth, H.C. (2003). Combining nonthermal technologies to control food borne microorganism. *Int J Food Microbiol* 89:125-38.
45. Sado, P.N, Jinneman, K.C, Husby, G.J, Sorg, S.M, Omiecinski, C.J. (1998). Identification of *L. monocytogenes* from unpasteurized apple juice using rapid test kits. *J Food Prot* 61:1199-202.
46. Splittstoesser, D.F. (1991). Fungi of importance in processed fruits. In: Arora DK, Mukerji KG, Marth EH, editors. Handbook of applied mycology. New York: Marcel Dekker, Inc. p 201-19.
47. Splittstoesser, D. F., and A. D. King, (1984). In M. Speck (ed.), "Compendium of methods for the microbiological

Isolation Of Heat Resistant Fungi From Canned Fruits

examination of foods," 2nd ed. American Public Health Association, Washington, D. C.

48. Stephanie, T. D. and Martha, B. S. (2008), Microsoft Encarta, 2008. Microsoft Corporation. U.S.A

49. Stolk, A.C. and Samson, R.A, (1981) studies on Talaromyces and related genera. Persoonia 6, 341-357.

50. Uzuegbu J.O. and A. T. Emifoniye, (1984). Post harvest fungal spoilage of some Nigerian fruits and vegetables. Journal of Nigerian food. Pp153-155.

51. Walker W, Phillips CA. (2008). The effect of preservatives on Alicyclobacillus acidoterrestris and Propionibacterium cyclohexanicum in fruit juice. Food Contr 19:974—81.

52. Wheeler, B. E. (1982). An introduction to plant disease. John Wiley and sons' Ltd. Chichester. Pp283-299



Sani Ibrahim

Assistant Professor, Department of Biotechnology, Faculty of Science and Humanities, SRM University, Kattankulathur.

Publish Research Article International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication, you will be pleased to know that our journals are

Associated and Indexed, India

- * International Scientific Journal Consortium
- * OPEN J-GATE

Associated and Indexed, USA

- EBSCO
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Database
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Golden Research Thoughts
258/34 Raviwar Peth Solapur-413005, Maharashtra
Contact-9595359435
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com
Website : www.aygrt.isrj.net