

---

# International Multidisciplinary Research Journal

## *Golden Research Thoughts*

Chief Editor  
Dr.Tukaram Narayan Shinde

---

Publisher  
Mrs.Laxmi Ashok Yakkaldevi

Associate Editor  
Dr.Rajani Dalvi

Honorary  
Mr.Ashok Yakkaldevi

## Welcome to GRT

**RNI MAHMUL/2011/38595**

**ISSN No.2231-5063**

Golden Research Thoughts Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

## Regional Editor

Manichander Thammishetty  
Ph.d Research Scholar, Faculty of Education IASE, Osmania University, Hyderabad

## International Advisory Board

Kamani Perera Regional Center For Strategic Studies, Sri Lanka	Mohammad Hailat Dept. of Mathematical Sciences, University of South Carolina Aiken	Hasan Baktir English Language and Literature Department, Kayseri
Janaki Sinnasamy Librarian, University of Malaya	Abdullah Sabbagh Engineering Studies, Sydney	Ghayoor Abbas Chotana Dept of Chemistry, Lahore University of Management Sciences[PK]
Romona Mihaila Spiru Haret University, Romania	Ecaterina Patrascu Spiru Haret University, Bucharest	Anna Maria Constantinovici AL. I. Cuza University, Romania
Delia Serbescu Spiru Haret University, Bucharest, Romania	Loredana Bosca Spiru Haret University, Romania	Ilie Pintea, Spiru Haret University, Romania
Anurag Misra DBS College, Kanpur	Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Xiaohua Yang PhD, USA
Titus PopPhD, Partium Christian University, Oradea,Romania	George - Calin SERITAN Faculty of Philosophy and Socio-Political Sciences AL. I. Cuza University, Iasi	.....More

## Editorial Board

Pratap Vyamktrao Naikwade ASP College Devrukh,Ratnagiri,MS India	Iresh Swami Ex - VC. Solapur University, Solapur	Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur
R. R. Patil Head Geology Department Solapur University,Solapur	N.S. Dhaygude Ex. Prin. Dayanand College, Solapur	R. R. Yaliker Director Managment Institute, Solapur
Rama Bhosale Prin. and Jt. Director Higher Education, Panvel	Narendra Kadu Jt. Director Higher Education, Pune	Umesh Rajderkar Head Humanities & Social Science YCMOU,Nashik
Salve R. N. Department of Sociology, Shivaji University,Kolhapur	K. M. Bhandarkar Praful Patel College of Education, Gondia	S. R. Pandya Head Education Dept. Mumbai University, Mumbai
Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai	Sonal Singh Vikram University, Ujjain	Alka Darshan Shrivastava Shaskiya Snatkottar Mahavidyalaya, Dhar
Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College, Indapur, Pune	G. P. Patankar S. D. M. Degree College, Honavar, Karnataka	Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore
Awadhesh Kumar Shirotriya Secretary,Play India Play,Meerut(U.P.)	Maj. S. Bakhtiar Choudhary Director,Hyderabad AP India.	S.KANNAN Annamalai University,TN
	S.Parvathi Devi Ph.D.-University of Allahabad	Satish Kumar Kalhotra Maulana Azad National Urdu University
	Sonal Singh, Vikram University, Ujjain	

Address:-Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India  
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.aygrt.isrj.org

# Golden Research Thoughts

International Recognition Multidisciplinary Research Journal

ISSN: 2231-5063

Impact Factor : 3.4052(UIF)

Volume - 5 | Issue - 9 | March - 2016



## NAKSHATRA BASED RAINFALL ANALYSIS AND ITS IMPACT ON CROPS DURING MONSOON SEASON AT GKVK STATION



**K. N. Krishnamurthy<sup>1</sup>, D. M. Gowda<sup>2</sup> And T. Vasanthkumar<sup>3</sup>**

<sup>1</sup>Associate professor, Dept. of Statistics, UAS, GKVK, Bengaluru.

<sup>2</sup>Professor, Dept. of Statistics, UAS, GKVK, Bengaluru.

<sup>3</sup>Assistant professor, Dept. of Statistics, UAHS, Mudigere.

---

### ABSTRACT:

Agricultural creation in India principally relies on the storm precipitation. Its auspicious onset, spread over the nation and appropriation is vital for Kharif crops. One of the conventional ways the ranchers of India when all is said in done and its southern States specifically examine precipitation examples is through sun powered groups of stars called "Nakshatras". In the present paper, a brief investigation of precipitation taking into account Nakshatra periods is considered. The every day precipitation information for a time of 30 years from 1984 - 2013 was gathered from the agrometeorological focus situated at GKVK, Bengaluru. The examination uncovers that the periods from Punarvasu to Chitta which covers the storm period got great measure of precipitation amid which trims like groundnut, sunflower, maize, finger millet and heartbeats like redgram, pigeon pea and chick pea can be taken up. Further, the pattern in the vast majority of the Nakshatra periods was diminishing aside from Makha and Chitta. This demonstrates throughout the years the aggregate precipitation got

---

amid these periods were diminishing. There is a need to draw up an emergency course of action to moderate the best possible arranging of rainfed yields by agriculturists.

**KEY WORDS:** Nakshatra-wise rainfall, Constellation, Trend analysis of rainfall

#### **INTRODUCTION:**

Agriculture is the backbone of Indian economy. Agricultural production mainly depends upon the occurrence of rainfall during the cropping season. The timely onset, its distribution and sufficient monsoon rainfall is the key for better agricultural production in the country which directly influences rural poverty alleviation (Varshneya et al, 2011). There is considerable traditional knowledge of variability of rainfall patterns, since rainfed cultivation has been carried out for several centuries in India. The periods used by the farmers are however, not weeks or months but so-called "Nakshatras" which are 13- or 14-day periods based on calendar. The Nakshatras are constellations through which the sun passes in a year. There are 27 Nakshatras in a year viz., Purvashada, Uttarabhadrapada, Shravana, Danista, Purvabhadra, Uttarabhadra, Revathi, Ashwini, Bharini, Krutika, Rohini, Mrugashira, Aridhra, Punarvasu, Pushya, Aslesha, Makha, Pubba, Uttara, Hastha, Chitta, Swathi, Vishaka, Anuradha, Jyeshtha and Moola Nakshatras. Of these, the periods from Rohini to Chitta Nakshatras cover the monsoon season. The Nakshatra commences when the sun enters the specific constellation. Thus, the knowledge of the variability in these time units rather than weeks or months is considered important by the traditional farmers in Karnataka and other neighboring States. The appropriate time for farming operations can also be worked out in terms of these time periods (Subash et al., 2011). In order to translate the meteorological events into farmer's terminology, it is necessary to perform rainfall analysis in Nakshatra periods. Not many reviews are available but however, the following recent studies have been documented.

Narahari Rao et al. (2000) have suggested a crop model PNUTGRO based on Nakshatra periods in which an optimum sowing window for rainfed groundnut crop for Ananthpur region has been given. Gadgil and Rao (2000) have developed an approach for identifying appropriate strategies for rainfed regions based primarily on information and prediction of climate variability during Nakshatra periods. De et al. (2004) performed a time series analysis of rainfall on different Nakshatra periods covering Indian monsoon season. Bavadekar et al. (2008) have carried out Nakshatra-wise rainfall analysis for drought prone areas of Maharashtra. Chinnchorkar et al. (2013) on their study of rainfall variability at Anand in middle Gujarat region have concluded that rainfall during Punarvasu and Pushya is more assured than in other Nakshatras, while it is least assured during Makha and Pubba Nakshatras. Chhabra and Haris (2014) have compiled the indigenous knowledge related to climatic parameters, their forecasting during different time periods of a year (Nakshatras) based on experiences of the farmers and comparing indigenous knowledge with modern scientific analysis of weather data and their relationship with wheat and rabi maize yield in Patna, Bihar. Hazra et al. (2014) have proposed a mixture model of two distributions for each Nakshatra periods for the Eastern plateau of India. In the present study Nakshatra based rainfall analysis has been carried out in order to study its trend in each Nakshatra period as well as to study its impact on the crops during monsoon at GKVK station.

#### **METHODOLOGY**

The daily rainfall data recorded at Agro meteorology centre, UAS, GKVK, Bengaluru for 30 years from 1984 to 2013 was used to analyze Nakshatra-wise rainfall distribution at GKVK station. Of the 27 Nakshatras, 11 Nakshatras from Rohini (May 25 to June 7) to Chitta (Oct.11-Oct.23) were considered for

the analysis. The mean, standard deviation, coefficient of variation (CV%), minimum and maximum for Nakshatra-wise rainfall were calculated. The trend analysis of rainfall at each Nakshatra period has been carried out.

## RESULTS AND DISCUSSION

### Characterization of Nakshatra-wise rainfall

Eleven Nakshatras were considered for analysis because this period coincides with the crop growing period of Kharif season. The descriptive statistics of Nakshatra-wise rainfall is presented in Table.1. Rainfall was received in all Nakshatras and good amount was received from Aslesha (Aug.3-Aug.16) to Chitta (Oct.11-Oct.23). The maximum total rainfall of 370.8 mm occurred during Aslesha followed by 302.5 mm during Uttara. The mean rainfall was maximum during Hastha (102.8 mm) followed by Uttara (99.8 mm). The least rainfall was received during Aridhra (21.2 mm). The CV of rainfall was lowest during Hastha (65.9 per cent) followed by Pushya (72.9 per cent) while it was highest (125.8 per cent) during Mrugashira. The pre monsoon mean rainfall was highest during Rohini (64.3 mm) and lowest during Mrugashira (30.6mm). On the other hand, during post monsoon season the highest was in Hastha (102.8 mm) and the least during Chitta (83.7 mm).

### Crop selection for rainfall in different Nakshatra periods

The Nakshatra-wise mean rainfall is shown in Fig.1. The figure shows that during Rohini Nakshatra a good amount of rainfall was received. This period can be utilized for sowing groundnut and red gram. Mrugashira and Aridhra Nakshatras received less rainfall during which one can take up crops like finger millet, sunflower, pigeon pea and groundnut. The period Rohini to Uttara received good rainfall which is suitable for finger millet, sunflower, maize, and pulses like red gram, pigeon pea, chick pea.

**Table 1: Descriptive Statistics for Nakshatra wise Rainfall (mm) at GVKK Station from 1984-2013**

Season	Nakshatra	Period	Mean	SD	CV(%)	Minimum	Maximum
Pre-monsoon	Rohini	May.25-Jun.7	64.3	49.4	76.8	0.0	163.3
	Mrugashira	Jun.8-Jun.21	30.6	38.5	125.8	0.0	149.6
Monsoon	Aridhra	Jun.22-Jul.5	21.2	16.4	77.4	0.0	58.0
	Punarvasu	Jul.6-Jul.19	48.5	42.5	87.6	0.8	183.2
	Pushya	Jul.20-Aug.2	53.9	39.3	72.9	9.6	154.4
	Aslesha	Aug.3-Aug.16	67.4	72.6	107.7	6.0	370.8
	Makha	Aug.17-Aug.30	68.1	58.8	86.3	1.4	204.2
	Pubba	Aug.31-Sep.12	60.4	51.3	84.9	3.6	170.0
	Uttara	Sep.13-Sep.26	99.8	84.6	84.8	0.0	302.6
Post-monsoon	Hastha	Sep.27-Oct.10	102.8	67.7	65.9	6.4	275.4
	Chitta	Oct.11-Oct.23	83.7	79.5	95.0	0.0	286.8

### Highest rainfall and Trend analysis of rainfall during Nakshatra periods

The highest rainfall during Nakshatra periods are presented in Table 2. The rainfall during Nakshatra periods ranged from 58.0 mm to 370.8 mm. Aslesha Nakshatra received the maximum rainfall of 370.8 mm during 1997 followed by Uttara Nakshatra of 302.6 mm. The low rainfall of 58.0

mm was recorded for Aridhra during 1992. The trend equations fitted for each of the Nakshatra periods reveal that all of them showed decreasing trend over the time period except for Makha and Chitta Nakshatras where the trend showed an increasing pattern. The fitted line and its equation are depicted in Fig. 2.





Fig.1: Nakshatra-wise mean rainfall (mm) at GVKK station from 1984-2013.

**Table 2: Nakshatra-wise highest Rainfall (mm) and its trend equations**

Season	Nakshatra	Period	Highest Rainfall		Trend equation
			Amount (mm)	Year	
Pre-monsoon	Rohini	May.25-Jun.7	163.3	1991	$Y = 1405.0 - 0.670X$
	Mrugashira	Jun.8-Jun.21	149.6	1996	$Y = 608.8 - 0.289X$
Monsoon	Aridhra	Jun.22-Jul.5	58.0	1992	$Y = 362.5 - 0.170X$
	Punarvasu	Jul.6-Jul.19	183.2	1988	$Y = 457.6 - 0.204X$
	Pushya	Jul.20-Aug.2	154.4	2007	$Y = 115.9 - 0.031X$
	Aslesha	Aug.3-Aug.16	370.8	1997	$Y = 471.0 - 0.202X$
	Makha	Aug.17-Aug.30	204.2	2008	$Y = -3688.0 + 1.879X$
	Pubba	Aug.31-Sep.12	170.0	1993	$Y = 1265.0 - 0.603X$
	Uttara	Sep.13-Sep.26	302.6	2001	$Y = 3912.0 - 1.907X$
Post-monsoon	Hastha	Sep.27-Oct.10	275.4	1997	$Y = 686.3 - 3.382X$
	Chitta	Oct.11-Oct.23	286.8	1999	$Y = -135.5 + 0.109X$

The pre monsoon period like Rohini, received an average of 64.3 mm rainfall during which land preparation and sowing of crops can be taken up while the period from Punarvasu to Chitta Nakshatras are suited for the sowing of the crops that are dependent on monsoon.









Fig.2: Nakshatra-wise Trend analysis of rainfall for the period from 1984 to 2013.







Fig.2: Nakshatra-wise Trend analysis of rainfall for the period from 1984 to 2013(Contd.)








Fig.2: Nakshatra-wise Trend analysis of rainfall for the period from 1984 to 2013 (Contd.)

### CONCLUSION

From the above examination it uncovers that the periods from Punarvasu to Chitta which covers the rainstorm period got great measure of precipitation amid which trims like groundnut, sunflower, maize, finger millet and heartbeats like redgram, pigeon pea and chick pea can be taken up. Further, in the vast majority of the Nakshatra periods the pattern was diminishing with the exception of Makha and Chitta. This demonstrates throughout the years the aggregate precipitation got amid these periods were diminishing. Hence, there is a need to draw up contingent planning of rainfed crops by farmers during these periods.

### REFERENCES

1. BAVADEKAR, V. R., JADHAV, J. D., MOKASHI, D.D., KHADTARE, S.V., AND KADAM, J. R. 2008. Rainfall variation and probabilities in different Nakshatras of drought prone areas, *Ag. Update*, 3(1&2): 153-158.
2. CHABBRA, V., AND HARIS, A. A., 2014. Nakshatra based rainfall analysis and its impact on rabi crops yield for Patna, Bihar, *Sch. J. Agric. Vet. Sci.*, 1(4): 168-172.
3. CHINCHORKAR, S.S., VAIDYA, V. B., VYAS PANDEY., AND DAMLE, K.S. 2013. Nakshatra-wise Rainfall variability at Anand in Middle Gujarat Region, *Asian Agri.-History*, 17(4): 389-393.
4. DE, U. S., JOSHI, U.R., AND PRAKASH RAO, G. S., 2004. Nakshatra based rainfall climatology, *Mausam*, 55(2): 305-312.
5. GADGIL SULOCHANA., AND RAO SESHAGIRI, P. R., 2000. Farming strategies for a variable climate – A challenge, *Current Science*, 78(10): 1203-1215.
6. HAZRA ARNAB., BHATTACHARYA, S., AND BANIK, P., 2014. Modelling Nakshatra-wise rainfall data of the Eastern plateau region of India, *Mausam*, 65(2): 264-270.

7. NARAHARI RAO, K., SULOCHANA GADGIL., SESHAGIRI RAO, P. R., AND SAVITHRI, K., 2000. Tailoring strategies to rainfall variability – The choice of the sowing window, *Current Science*, 78(10): 1216-1230.
8. SUBASH, N., SINGH, S. S., AND NEHA PRIYA., 2011. Nakshatra based rainfall variability, trends and its influences on rice wheat production- A case study over two cities in Bihar, India. *Journal of Agrometeorology*, 13(1): 31-37.
9. VARSHNEYA, M. C., VAIDYA, V. B., NANAJI KALE., AND KETAN KALE., 2011. Performance and of Sakshat Krishi project in India, *Asian Agri.- History*, 14(4): 361-372.



Associate professor, Dept. of Statistics, UAS, GKVK, Bengaluru.

# Publish Research Article International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication, you will be pleased to know that our journals are

## Associated and Indexed, India

- \* International Scientific Journal Consortium
- \* OPEN J-GATE

## Associated and Indexed, USA

- EBSCO
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Database
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Golden Research Thoughts  
258/34 Raviwar Peth Solapur-413005, Maharashtra  
Contact-9595359435  
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com  
Website : www.aygrt.isrj.org