



# Golden Research Thoughts

GRT



## THE MINIMUM GLOBAL DOMINATING ENERGY OF A GRAPH

K.B.Murthy<sup>1</sup>, Puttaswamy<sup>2</sup>

<sup>1</sup>Department of Mathematics, College of Agriculture, University of Agricultural Sciences, GKVK, Bangalore- 560065, India.

<sup>2</sup>Department of Mathematics, P.E.S.College of Engineering, Mandya, India.

### Abstract


In a graph  $G(V, E)$ , a subset  $D \subseteq V$  is called a global dominating set of  $G$ , if  $D$  is a dominating set of both  $G$  and  $\bar{G}$ . The global domination number  $\gamma_g(G)$  is the minimum cardinality of a minimal global dominating set in  $G$ . In this paper, we study the minimum dominating energy, denoted by  $E_{GD}(G)$ , of a graph  $G$  and minimum double dominating energy of  $G$ . We compute the minimum global dominating energies of complete graph, complete bipartite graph, star graph and cocktail party graph. Upper and lower bounds for  $E_{GD}(G)$  are established.

### Mathematics Subject

Classification: 05C50, 05C69.

### Key words:

global dominating set, global domination number, global dominating matrix, global dominating eigenvalues, minimum global dominating energy, Minimum Double Dominating Energy.



### 1. INTRODUCTION

In this paper, by a graph  $G(V, E)$  we mean a simple graph that is finite, have no loops no multiple and directed edges. We denoted by  $n$  and  $m$  to the number of its vertices and edges, respectively. We refer the reader to Harary book [9] for more graph theoretical analogist not

defined here. A subset  $D \subseteq V(G)$  is called a dominating set of  $G$  if every vertex  $v \in V - D$  is adjacent to some vertex in  $D$ . The domination number  $\gamma(G)$  of  $G$  is the minimum cardinality of a minimal dominating set in  $G$ . A subset  $D \subseteq V(G)$  is called a global dominating set in  $G$  if  $D$  is a dominating set of both  $G$  and  $\bar{G}$ . The global domination number  $\lambda_g(G)$  is the minimum cardinality of a minimal global dominating set in  $G$ . Any global dominating set in a graph with minimum cardinality is called a minimum global dominating set. The concept of global domination in graph was introduced by Sampathkumar [18], while Kulli and Janakiram [14] have introduced the concept of total global dominating sets. Also, there are more studies and details in this concept can be see it in [4, 20, 21, 19] and the references cited there in. While for more details in domination theory of graphs we refer to Haynes et al. book [10].

The concept of energy of a graph was introduced by I. Gutman [7] in the year 1978. Let  $G$  be a graph with  $n$  vertices and  $m$  edges and let  $A = (a_{ij})$  be the adjacency matrix of the graph. The eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  of  $A$ , assumed in non increasing order, are the eigenvalues of the graph  $G$ . As  $A$  is real symmetric, the eigenvalues of  $G$  are real with sum equal to zero. The energy  $E(G)$  of  $G$  is defined to be the sum of the absolute values of the eigenvalues of  $G$ , i.e.

$$E(G) = \sum_{i=1}^n |\lambda_i|.$$

For more details on the mathematical aspects of the theory of graph energy see [2, 8, 15]. The basic properties including various upper and lower bounds for energy of a graph have been established in [16, 17], and it has found remarkable chemical applications in the molecular orbital theory of conjugated molecules [5, 6].

Recently C. Adiga et al [1] defined the minimum covering energy,  $E_C(G)$  of a graph which depends on its particular minimum cover  $C$ . Further, minimum dominating energy, Laplacian minimum dominating energy and minimum dominating distance energy of a graph  $G$  can be found in [11, 12, 13] and the references cited there in.

Motivated by these papers, we study the minimum global dominating energy  $E_{GD}(G)$  of a graph  $G$  and minimum double dominating Energy. We compute minimum global dominating energies of some standard graphs. Upper and lower bounds for  $E_{GD}(G)$  are established. It is possible that the upper dominating energy that we are considering in this paper may be have some applications in chemistry as well as in other areas.

## 2. THE MINIMUM GLOBAL DOMINATING ENERGY OF A GRAPH

Let  $G$  be a graph of order  $n$  with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(G)$ . Let  $D$  be a global dominating set in  $G$ . The global domination number  $\gamma_g(G)$  of  $G$  is the cardinality of a smallest global dominating set in  $G$ . Any global dominating set  $D$  with cardinality equals to  $\gamma_g(G)$  is called a minimum global dominating set of  $G$ . The minimum global dominating matrix of  $G$  is then  $n \times n$  matrix, denoted by  $AGD(G) = (a_{ij})$ , where Characteristic polynomial of  $AGD(G)$  is denoted by

$$f_n(G, \lambda) = \det(\lambda I - A_{GD}(G)).$$

The minimum global dominating eigen values of a graph  $G$  are the eigenvalues of  $A_{GD}(G)$ . Since  $A_{GD}(G)$  is real and symmetric, its eigen values are real numbers and we label them in non-increasing order  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ . The minimum global dominating energy of  $G$  is defined as:

$$EGD(G) = \sum_{i=1}^n |\lambda_i|.$$

We first compute the minimum global dominating energy of a graph in Figure 1




Figure-1

Let  $G$  be a graph in Figure 1, with vertices set  $\{v_1, v_2, v_3, v_4, v_5, v_6\}$ . Then  $G$  has more one set as a minimum global dominating set. For example,  $D_1 = \{v_1, v_3, v_4\}$ . Then

$$A_{GD_1}(G) = \begin{pmatrix} 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 1 & 0 \end{pmatrix}$$

The characteristic polynomial of  $A_{GD_1}(G)$  is  $f_n(G, \lambda) = \lambda^6 - 3\lambda^5 - 5\lambda^4 + 12\lambda^3 + 9\lambda^2 - 9\lambda - 5$ .

Hence, the minimum global dominating eigenvalues are  $\lambda_1 \approx 3.2263$ ,  $\lambda_2 \approx 1.9102$ ,  $\lambda_3 \approx 1.0000$ ,  $\lambda_4 \approx -0.4939$ ,  $\lambda_5 \approx -1$ ,  $\lambda_6 \approx -1.6426$ .

Therefore the minimum global dominating energy of  $G$  is  $EGD_1(G) \approx 9.2730$ .

But if we take another minimum global dominating set of  $G$ , namely  $D_2 = \{v_2, v_3, v_6\}$ , we get that

$$A_{GD_2}(G) = \begin{pmatrix} 0 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 1 & 1 \end{pmatrix}$$

The characteristic polynomial of  $A_{GD_2}(G)$  is  $f_n(G, \lambda) = \lambda^6 - 3\lambda^5 - 5\lambda^4 + 11\lambda^3 + 8\lambda^2 - 8\lambda - 4$ .

The minimum global dominating eigenvalues are  $\lambda_1 \approx 3.3795$ ,  $\lambda_2 \approx 1.6708$ ,  $\lambda_3 \approx 1$ ,  $\lambda_4 \approx -0.4399$ ,  $\lambda_5 \approx -1$ ,  $\lambda_6 \approx -1.6105$ . Therefore the minimum global dominating energy of  $G$  is  $EGD_2(G) \approx 9.1007$ .

This example illustrates the fact that the minimum global dominating energy of a graph  $G$  depends on the choice of the minimum global dominating set. i.e. the minimum global dominating energy is not a graph invariant.

In the following section, we introduce some properties of characteristic polynomial of minimum global dominating matrix of a graph  $G$ .

**Theorem 2.1.** Let  $G$  be a graph of order  $n$ , size  $m$ , minimum global dominating set  $D$  and let  $f_n(G, \lambda) = c_0\lambda^n + c_1\lambda^{n-1} + c_2\lambda^{n-2} + \dots + c_n$ .

be the characteristic polynomial of the minimum global dominating matrix of a graph  $G$ .

Then

- 1  $c_0 = 1$ .
- 2  $c_1 = -|D|$
3.  $c_2 = \binom{|D|}{2} - 2$ .

**Proof.** 1. From the definition of  $f_n(G, \lambda)$ .

2. Since the sum of diagonal elements of  $AGD(G)$  is equal to  $|D|$ . The sum of determinants of all  $1 \times 1$  principal submatrices of  $AGD(G)$  is the trace of  $AGD(G)$ , which evidently is equal to  $|D|$ . Thus,  $(-1)^1 c_1 = |D|$ .

3.  $(-1)^2 c_2$  is equal to the sum of determinants of all  $2 \times 2$  principal submatrices of  $AGD(G)$ , that is

$$\begin{aligned} c_2 &= \sum_{1 \leq i < j \leq n} \begin{vmatrix} a_{ii} & a_{ij} \\ a_{ji} & a_{jj} \end{vmatrix} \\ &= \sum_{1 \leq i < j \leq n} (a_{ii}a_{jj} - a_{ij}a_{ji}) \\ &= \sum_{1 \leq i < j \leq n} a_{ii}a_{jj} - \sum_{1 \leq i < j \leq n} a_{ij}^2 \\ &= \binom{|D|}{2} - m \end{aligned}$$

**Theorem 2.2.** Let  $G$  be a graph of order  $n$ . Let  $\lambda_1, \lambda_2, \dots, \lambda_n$  be the eigenvalues of

$AGD(G)$ . Then

- (i)  $\sum_i^n \lambda_i = |D|$ .
- (ii)  $\sum_i^n \lambda_i^2 = |D| + 2m$ .

Proof. (i) Since the sum of the eigen values of  $AGD(G)$  is the trace of  $AGD(G)$ , then

$$\sum_i^n \lambda_i = \sum_{i=1}^n a_{ii} = |D|.$$

(ii) Similarly the sum of squares eigenvalues of  $AGD(G)$  is the trace of  $(AGD(G))^2$ .

Then,

$$\begin{aligned} \sum_{i=1}^n \lambda_i^2 &= \sum_{i=1}^n \sum_{j=1}^n a_{ij} a_{ji} \\ &= \sum_{i=1}^n a_{ii}^2 + \sum_{i \neq j} a_{ij} a_{ji} \\ &= \sum_{i=1}^n a_{ii}^2 + 2 \sum_{i \neq j} a_{ij}^2 \\ &= |D| + 2m \end{aligned}$$

Bapat and S.Pati [3], proved that if the graph energy is a rational number then it is an even integer. Similar result for minimum global dominating energy is given in the following theorem.

**Theorem 2.3.** Let  $G$  be a graph with a minimum global dominating set  $D$ . If the minimum global dominating energy  $EGD(G)$  of  $G$  is a rational number, then

$$EGD(G) \equiv |D| \pmod{2}.$$

**Proof.** Let  $\lambda_1, \lambda_2, \dots, \lambda_n$  be the minimum global dominating eigen values of a graph  $G$  of which  $\lambda_1, \lambda_2, \dots, \lambda_r$  are positive and the rest are non-positive, then

$$\begin{aligned} \sum_{i=1}^n |\lambda_i| &= (\lambda_1 + \lambda_2 + \dots + \lambda_r) - (\lambda_{r+1} + \lambda_{r+2} + \dots + \lambda_n) \\ &= 2(\lambda_1 + \lambda_2 + \dots + \lambda_r) - (\lambda_1 + \lambda_2 + \dots + \lambda_r). \\ &= 2q - |D|. \text{ where } q = \lambda_1 + \lambda_2 + \dots + \lambda_r. \end{aligned}$$

Since  $\lambda_1, \lambda_2, \dots, \lambda_r$  are algebraic integers, so is their sum. Hence  $(\lambda_1 + \lambda_2 + \dots + \lambda_r)$  must be an integer if  $EGD(G)$  is rational.

**Theorem 2.4.** For  $n \geq 1$ , the minimum global dominating energy of the complete graph  $K_n$  is  $EGD(K_n) = n$ .

**Proof.** For the complete graphs  $K_n$ , the minimum global dominating set is all vertices, i.e.  $D = V(K_n)$ .

Then,

$$\begin{aligned} A_{GD}(K_n) &= \begin{pmatrix} 1 & 1 & \dots & 1 \\ 1 & 1 & \dots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \dots & 1 \end{pmatrix}_{n \times n} \\ &= \lambda^{n-1}(\lambda - n) \end{aligned}$$

The characteristic polynomial of  $AGD(K_n)$  is

$$f_n(K_n, \lambda) = \begin{vmatrix} \lambda - 1 & -1 & \cdots & -1 \\ -1 & \lambda - 1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & \lambda - 1 \end{vmatrix}_{n \times n}$$

The spectrum of  $K_n$  is

$$GD\ Spec(K_n) = \begin{pmatrix} 0 & n \\ n - 1 & 1 \end{pmatrix}$$

Therefore, the minimum global dominating energy of the complete graph is

$$EGD(K_n) = n.$$

**Theorem 2.5.** For the complete bipartite graph  $K_{r,r}$ , for  $r \geq 2$ , the minimum global dominating energy is equal to

$$(r + 1) + \sqrt{r^2 + 2r - 3}$$

**Proof.** For the complete bipartite graph  $K_{r,r}$  with vertex set  $V = (V_1, V_2)$  where  $V_1$  and  $V_2$  are the partite sets of its,  $V_1 = \{v_1, v_2, \dots, v_r\}$  and  $V_2 = \{u_1, u_2, \dots, u_r\}$ .

The minimum global dominating set is  $D = \{v_1, u_1\}$ . Then

$$A_{GD}(K_{r,r}) = \begin{pmatrix} 1 & 0 & \cdots & 0 & 1 & 1 & \cdots & 1 \\ 0 & 0 & \cdots & 0 & 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 & 1 & 0 & \cdots & 0 \\ 1 & 1 & \cdots & 1 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 & 0 & 0 & \cdots & 0 \end{pmatrix}_{2r \times 2r}$$

The characteristic polynomial of  $AGD(K_{r,r})$  is

$$f_n(K_{r,r}, \lambda) = \begin{vmatrix} \lambda - 1 & 0 & \cdots & 0 & -1 & -1 & \cdots & -1 \\ 0 & \lambda & \cdots & 0 & -1 & -1 & \cdots & -1 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda & -1 & -1 & \cdots & -1 \\ 1 & 1 & \cdots & 1 & \lambda & 1 & 0 & \cdots & 0 \\ -1 & -1 & \cdots & -1 & 0 & \lambda & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ -1 & -1 & \cdots & -1 & 0 & 0 & \cdots & \lambda \end{vmatrix}_{2r \times 2r}$$

$$= \lambda^{2r-4} [\lambda^2 + (r-1)\lambda - (r-1)] [\lambda^2 - (r+1)\lambda + (r-1)]$$

The spectrum of  $K_{r,r}$  is  $GD\text{Spec}(K_{r,r})$

$$\begin{pmatrix} 0 & \frac{(1-r)-\sqrt{r^2+2r-3}}{2} & \frac{(1-r)+\sqrt{r^2+2r-3}}{2} & \frac{(1+r)-\sqrt{r^2-2r+5}}{2} & \frac{(1+r)+\sqrt{r^2-2r+5}}{2} \\ 2r-4 & 1 & 1 & 1 & 1 \end{pmatrix}$$

Therefore, the minimum global dominating energy of the complete bipartite graph

is  $E_{GD}(K_{r,r}) = (1+r) + \sqrt{r^2+2r-3}$

**Theorem 2.6.** For  $n \geq 2$ , the minimum global dominating energy of a star graph

$K_{1,n-1}$  is less than or equal to  $2 + 2\sqrt{n-2}$

**Proof.** Let  $K_{1,n-1}$  be a star graph with vertex set  $V = \{v_0, v_1, v_2, \dots, v_{n-1}\}$ , where  $v_0$  is the central vertex. Since  $\overline{K_{n-1}} = K_1 \cup K_{n-1}$  it follows that the minimum global dominating set of  $K_{1,n-1}$  is  $D = \{v_0, v_i\}$  for any  $i = 1, 2, \dots, n-1$ .

Hence, if we choose  $D = \{v_0, v_i\}$ , then the minimum global dominating matrix of  $K_{1,n-1}$  is

$$A_{GD}(K_{1,n-1}) = \begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ 1 & 1 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & 0 & 0 & \dots & 0 \end{pmatrix}_{n \times n}$$

The characteristic polynomial of  $A_{GD}(K_{1,n-1})$  is

$$\begin{aligned} f_n(K_{1,n-1}, \lambda) &= \begin{vmatrix} \lambda-1 & -1 & -1 & \dots & -1 \\ -1 & \lambda-1 & 0 & \dots & 0 \\ -1 & 0 & \lambda & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -1 & 0 & 0 & \dots & \lambda \end{vmatrix}_{n \times n} \\ &= (\lambda)^{n-3} [\lambda^3 - 2\lambda^2 - (n-2)\lambda + (n-2)] \\ &\leq (\lambda)^{n-3} [\lambda^3 - 2\lambda^2 - (n-2)\lambda + 2(n-2)] \end{aligned}$$

The spectrum of  $K_{1,n-1}$  is

$$GD\text{Spec}(K_{1,n-1}) = \begin{pmatrix} 0 & 2 & \sqrt{n-2} & -\sqrt{n-2} \\ n-3 & 1 & 1 & 1 \end{pmatrix}$$

Therefore, the minimum global dominating energy of a star graph is

$$E_{GD}(K_{1,n-1}) \leq 2 + 2\sqrt{n-2}$$

**Definition 2.7.** The cocktail graph, denoted by  $K_{2 \times p}$ , is a graph having vertex set  $V(K_{2 \times p}) = \cup_{i=1}^p \{u_i, v_i\}$  and edge set  $E(K_{2 \times p}) = \{u_i u_j, v_i v_j, u_i v_j, v_i u_j : 1 \leq i < j \leq p\}$ , i.e,  $n = 2p, m = \frac{p^2-3p}{2}$  and for ever  $v \in V(K_{2 \times p}), d(v) = 2p - 2$ .

**Theorem 2.8.** For the cocktail party graph  $K_{2 \times p}$  of order  $2p$ , for  $p \geq 3$ , the minimum global dominating energy is equal to  $(2p - 1) + \sqrt{5}(p - 1)$

**Proof.** For cocktail party graphs  $K_{2 \times p}$  the minimum domination set is  $D = \{u_i | 1 \leq i \leq p\}$ . Hence, for cocktail party graphs the minimum global dominating matrix is

$$A_{GD}(K_{2 \times p}) = \begin{pmatrix} 1 & 0 & 1 & 1 & \cdots & 1 & 1 \\ 0 & 0 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & 1 & 0 & \cdots & 1 & 1 \\ 1 & 1 & 0 & 0 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & 1 & 1 & \cdots & 1 & 0 \\ 1 & 1 & 1 & 1 & \cdots & 0 & 0 \end{pmatrix}_{2p \times 2p}$$

The characteristic polynomial of  $A_{GD}(K_{2 \times p})$  is

$$f_n(K_{2 \times p}, \lambda) = \begin{vmatrix} 1 & 0 & 1 & 1 & \cdots & 1 & 1 \\ 0 & 0 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 1 & 1 & 0 & \cdots & 1 & 1 \\ 1 & 1 & 0 & 0 & \cdots & 1 & 1 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 1 & 1 & 1 & 1 & \cdots & 1 & 0 \\ 1 & 1 & 1 & 1 & \cdots & 0 & 0 \end{vmatrix}_{2p \times 2p}$$

$$= [\lambda^2 - (2p - 1)\lambda + (p - 1)][\lambda^2 + \lambda - 1]^{(p-1)}$$

The spectrum of  $K_{2 \times p}$  is

$$GD \text{ Spec}(K_{2 \times p}) = \left( \begin{array}{cccc} \frac{(2p-1) - \sqrt{4p^2 - 8p + 2}}{2} & \frac{(2p-1) + \sqrt{4p^2 - 8p + 2}}{2} & \frac{-1 - \sqrt{5}}{2} & \frac{-1 + \sqrt{5}}{2} \\ 1 & 1 & p - 1 & p - 1 \end{array} \right)$$

Therefore, the minimum global dominating energy

$$E_{GD}(K_{2 \times p}) = (2p - 1) + \sqrt{5}(p - 1).$$

**Theorem 2.9.** Let  $G$  be a graph of order  $n$  and size  $m$ . Then

$$\sqrt{2m + \gamma_g(G)} \leq E_{GD}(G) \leq \sqrt{n(2m + \gamma_g(G))}$$

Proof. Consider the Cauchy-Schwartz inequality

$$\left( \sum_{i=1}^n a_i b_i \right)^2 \leq \left( \sum_{i=1}^n a_i^2 \right) \left( \sum_{i=1}^n b_i^2 \right).$$

By choose  $a_i = 1$  and  $b_i = |\lambda_i|$ , we get

$$\begin{aligned} (E_{GD}(G))^2 &= \left( \sum_{i=1}^n |\lambda_i| \right)^2 \leq \left( \sum_{i=1}^n 1 \right) \left( \sum_{i=1}^n \lambda_i^2 \right) \\ &\leq n(2m + |D|) \\ &\leq n(2m + \gamma_g(G)). \end{aligned}$$

Therefore, the upper bound is holds. For the lower bound, since

$$\left( \sum_{i=1}^n |\lambda_i| \right)^2 \geq \sum_{i=1}^n \lambda_i^2.$$

Then,  $E_{GD}(G)^2 \geq \sum_{i=1}^n \lambda_i^2 = 2m + |D| = 2m + \gamma_g(G)$

Therefore,  $E_{GD}(G) \geq \sqrt{2m + \gamma_g(G)}$ .

Similar to McClellands [17] bounds for energy of a graph, bounds for  $E_{GD}(G)$  are given in the following theorem.

**Theorem 2.10.** Let  $G$  be a graph of order  $n$  and size  $m$ , respectively.

If  $P = \det(A_{GD}(G))$ , then

$$E_{GD}(G) \geq \sqrt{2m + \gamma_g(G) + n(n-1)P^{\frac{2}{n}}}$$

$$(E_{GD}(G))^2 = \left( \sum_{i=1}^n |\lambda_i| \right)^2 = \left( \sum_{i=1}^n |\lambda_i| \right) \left( \sum_{i=1}^n |\lambda_i| \right) = \sum_{i=1}^n |\lambda_i|^2 + 2 \sum_{i \neq j} |\lambda_i| |\lambda_j|.$$

$$\frac{1}{n(n-1)} \sum_{i \neq j} |\lambda_i| |\lambda_j| \geq \left( \prod_{i \neq j} |\lambda_i| |\lambda_j| \right)^{\frac{1}{n(n-1)}}.$$

$$\begin{aligned}
 (E_{GD}(G))^2 &\geq \sum_{i=1}^n |\lambda_i|^2 + n(n-1) \left( \prod_{i \neq j} |\lambda_i| |\lambda_j| \right)^{1/[n(n-1)]} \\
 &\geq \sum_{i=1}^n |\lambda_i|^2 + n(n-1) \left( \prod_{i=2}^n |\lambda_i|^{2(n-1)} \right)^{1/[n(n-1)]} \\
 &= \sum_{i=1}^n |\lambda_i|^2 + n(n-1) \left| \prod_{i \neq j} \lambda_i \right|^{2/n} \\
 &= 2u + \gamma_g(G) + u(n-1)P^{2/n}.
 \end{aligned}$$

**3.MINIMUM DOUBLE DOMINATING ENERGY:**

Let  $G$  be a simple graph of order  $n$  with vertex set  $V = \{v_1, v_2, v_3, \dots, v_n\}$  and edge set  $E$ . A subset  $D' \subseteq V$  is a double dominating set if  $D'$  is a dominating set and every vertex of  $V - D'$  is adjacent to atleast two vertices in  $D'$ . The double Domination number  $\gamma_{x2}(G)$  is the minimum cardinality taken over all the minimal double dominating sets of  $G$ .

Let  $D'$  be the minimum double dominating set of a graph. The minimum double dominating matrix of  $G$  is the  $n \times n$  matrix defined by  $A_D(G) = (a_{ij})$  where

$$a_{ij} = \begin{cases} 1 & \text{if } v_i v_j \in E \\ 1 & \text{if } i = j \text{ and } v_i \in D' \\ 0 & \text{otherwise} \end{cases}$$

The characteristic polynomial of  $A_D(G)$  is denoted by  $f_n(G, \lambda) = \det(\lambda I - A_D(G))$ .

The minimum double dominating eigen values of the graph  $G$  are the eigen values of  $A_D(G)$ .

Since  $A_D(G)$  is real and symmetric, its eigen values are real numbers and are labeled in non-increasing order  $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_n$ . the minimum double dominating energy of  $G$  is defined as  $E_D(G) = \sum_{i=1}^n |\lambda_i|$ .

**Example 3.1.** Let  $G$  be a cycle  $C_4$  on 4 vertices  $u_1, u_2, u_3, u_4$  with minimum double dominating set  $D' = \{u_1, u_3\}$ . Then

$$A_D(C_4) = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

The Characteristic polynomial of  $A_D(C_4)$  is  $\lambda^4 - 2\lambda^3 - 3\lambda^2 + 4\lambda$ , the minimum double dominating eigen values are  $0, 1, \frac{1+\sqrt{7}}{2}, \frac{1-\sqrt{7}}{2}$ , and the minimum double dominating energy is  $E_D(C_4) = 1 + \sqrt{17}$ .

**Theorem 3.2.** Let  $G$  be a graph with  $n$  vertices and  $m$  edges.

If  $\lambda_1, \lambda_2, \lambda_3, \dots, \lambda_n$  are the eigen values of  $A_D(G)$ , then  $\sum_{i=1}^n \lambda_i^2 = 2|E| + |D'|$ .

**Proof:** The sum of the squares of the eigen values of  $A_D(G)$  is the trace of  $A_D(G)^2$ .

$$\begin{aligned} \sum_{i=1}^n \lambda_i^2 &= \sum_{i=1}^n \sum_{j=1}^n a_{ij} a_{ji} \\ &= 2 \sum_{i < j} (a_{ij})^2 + \sum_{i=1}^n (a_{ij})^2 \\ &= 2|E| + |D'| \\ &= 2m + |D'|. \end{aligned}$$

**Theorem 3.3.** Let  $G$  be a simple graph with  $n$  vertices,  $m$  edges and let  $D'$  be a double dominating set of  $G$  and  $F = |\det A_D(G)|$  then ,

$$\sqrt{2m + |D'| + n(n-1)f^{\frac{2}{n}}} \leq E_D(G) \leq \sqrt{n(2m + |D'|)}$$

**Proof:** Let  $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots, \geq \lambda_n$  be the eigen values of  $A_D(G)$  .

By Cauchy-Schwarz inequality,  $\sum_{i=1}^n (a_i b_i)^2 \leq \sum_{i=1}^n (a_i^2) (\sum_{i=1}^n (b_i^2))$ .

Let  $a_i = 1$ ,  $b_i = |\lambda_i|$ ,

$$E_D(G)^2 = \left( \sum_{i=1}^n |\lambda_i| \right)^2 \leq n \left( \sum_{i=1}^n |\lambda_i|^2 \right) = n \sum_{i=1}^n \lambda_i^2 = n(2m + |D'|)$$

$$E_D(G) \leq \sqrt{n(2m + |D'|)}$$

$$[E_D(G)]^2 = \left( \sum_{i=1}^n |\lambda_i| \right)^2 = \sum_{i=1}^n |\lambda_i|^2 + \sum_{i=j} |\lambda_i| |\lambda_j|$$

From the inequality between the arithmetic and geometric mean,

We obtain  $\frac{1}{n(n-1)} \sum_{i=j} |\lambda_i| |\lambda_j| \geq \left( \prod_{i=j} |\lambda_i| |\lambda_j| \right)^{\frac{1}{n(n-1)}}$

$$\begin{aligned} \sum_{i \neq j} |\lambda_i| |\lambda_j| &\geq n(n-1) \left[ \prod_{i=1}^n |\lambda_i|^{2(n-1)} \right]^{\frac{1}{n(n-1)}} \\ &\geq n(n-1) \left[ \prod_{i=1}^n |\lambda_i| \right]^{\frac{2}{n}} \\ &\geq n(n-1) \left| \prod_{i=1}^n |\lambda_i| \right|^{\frac{2}{n}} \\ &\geq n(n-1) |\det A_D(G)|^{\frac{2}{n}} \end{aligned}$$

$$\begin{aligned} \sum_{i \neq j} |\lambda_i| |\lambda_j| &\geq n(n-1)F_n^{\frac{2}{n}} \\ [E_D(G)]^2 &\geq \sum_{i=1}^n |\lambda_i|^2 + n(n-1)F_n^{\frac{2}{n}} \\ &\geq (2m + |D'|) + n(n-1)F_n^{\frac{2}{n}} \\ [E_D(G)] &\geq \sqrt{2m + |D'| + n(n-1)F_n^{\frac{2}{n}}} \end{aligned}$$

**Definition 3.4.** The crown graph  $S_n^o$  for an integer  $n \geq 2$  is the graph with vertex set  $v = \{u_1, u_2, \dots, u_n, v_1, v_2, v_3, \dots, v_n\}$  and the edge set  $\{u_i v_j : 1 \leq i, j \leq n, i \neq j\}$ .

**Theorem 3.5.** For any  $n \geq 2$ , the double dominating energy of the crown graph  $S_n^o$  is equal to  $2 + 2(n-3) + \sqrt{n^2 - 2n + 9} + \sqrt{n^2 + 2n - 7}$ .

**Proof:** The crown graph  $S_n^o$  with vertex set  $v = \{u_1, u_2, \dots, u_n, v_1, v_2, v_3, \dots, v_n\}$  the minimum double dominating set  $D' = \{u_1, u_2, v_1, v_2\}$ .

$$\text{Then } A_{D'}(S_n^o) = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 & 1 & 1 & \dots & 1 \\ 0 & 1 & 0 & \dots & 0 & 1 & 0 & 1 & \dots & 1 \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 & 0 & \dots & 1 \\ \dots & \dots \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 & 1 & \dots & 0 \\ 0 & 1 & 1 & \dots & 1 & 1 & 0 & 0 & \dots & 0 \\ 1 & 0 & 1 & \dots & 1 & 0 & 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 1 & 0 & 0 & 0 & \dots & 0 \\ \dots & \dots \\ 1 & 1 & 1 & \dots & 0 & 0 & 0 & 0 & \dots & 0 \end{bmatrix}$$

Characteristic polynomial is

$$\begin{bmatrix} \lambda-1 & 0 & 0 & \dots & 0 & 0 & -1 & -1 & \dots & -1 \\ 0 & \lambda-1 & 0 & \dots & 0 & -1 & 0 & -1 & \dots & -1 \\ 0 & 0 & \lambda & \dots & 0 & -1 & -1 & 0 & \dots & -1 \\ \dots & \dots \\ 0 & 0 & 0 & \dots & \lambda & -1 & -1 & -1 & \dots & 0 \\ 0 & -1 & -1 & \dots & -1 & \lambda-1 & 0 & 0 & \dots & 0 \\ -1 & 0 & -1 & \dots & -1 & 0 & \lambda-1 & 0 & \dots & 0 \\ -1 & -1 & 0 & \dots & -1 & 0 & 0 & \lambda & \dots & 0 \\ \dots & \dots \\ -1 & -1 & -1 & \dots & 0 & 0 & 0 & 0 & \dots & \lambda \end{bmatrix}$$

Characteristic equation is

$$\lambda(\lambda - 2)(\lambda + 1)^{n-3}(\lambda - 1)^{n-3}(\lambda^2 + (n - 3)\lambda - (2n - 4)(\lambda^2 - (n - 1)\lambda - 2)) = 0$$

Minimum double dominating eigen values are

$$\lambda = 0, \lambda = 2, \lambda = -1, (n - 3 \text{ times}), \lambda = 1(n - 3 \text{ times})$$

$$\lambda = \frac{(n-1) \pm \sqrt{n^2 - 2n + 9}}{2} \text{ (one time each)}$$

$$\lambda = \frac{(3-n) \pm \sqrt{n^2 + 2n - 7}}{2} \text{ (one time each)}$$

Minimum double dominating energy

$$E_D(S_n^o) = 2 - 2(n - 3) + \sqrt{n^2 - 2n + 9} + \sqrt{n^2 + 2n - 7}.$$

## REFERENCES:

- [1] C.Adiga, A.Bayad, I.Gutman and S.A.Srikant, *The minimum covering energy of a graph*, K.J.Sci.,**34**(2012),39-56.
- [2] R.B.Bapat, *Graphs and Matrices*, Hindustan Book Agency,2011.
- [3] R.B.Bapat, S.Pati, *Energy of a graph is never an odd integer*, Bul. Kerala Math. Asso.,1(2011),129-132.
- [4] R.C. Brigham and R.D. Dutton, *Factor domination in graphs*, Dis.Math., **86** (1990),127-136.
- [5] A.Graovac, I.Gutman and N.Trinajstić, *Topological Approach to the Chemistry of Conjugated Molecules*,Springer,Berlin,1977.
- [6] I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer,Berlin,1986.
- [7] I. Gutman, *The energy of a graph*, Ber. Math-Statist. Sect. Fors. Graz, 103(1978), 1-22.
- [8]I.Gutman,X.Li,J.Zhang, *Graph Energy*,(Ed-s: M.Dehmer, F.Em-mert), Streib.,Anal.Comp.Net.,From Biology to Linguistics,Wiley-VCH,Weinheim, (2009),145-174.
- [9] F.Harary, *Graph Theory*, Addison Wesley, Massachusetts,1969.
- [10]T.W.Haynes,S.T.Hedetniemi and P.J.Slater, *Fundamentals of Domination in Graphs*, MarcelDekker,Inc.,NewYork1998.
- [11]R.Kanna,B.N.Dharmendra,and G.Sridhara, *Laplacian minimum dominating energy of a graph*, Int. J.Pure Appl.Math., **89:4**(2013), 565-581.
- [12]R.Kanna, B.N.Dharmendra, and G.Sridhara, *Minimum dominating distance energy of a graph*,J.Ind.Math. Soci.,**20**(2014),19-29.
- [13]R.Kanna, B.N.Dharmendra and G.Sridhara, *Minimum dominating energy of a graph*, Int. J.Pure Appl.Math., **85:4**(2013), 707-718.
- [14]V.R.Kulli and B.Janakiram,*The total global domination number of a graph*, Indian. J.Pure Appl.Math., **27**,(1996),537-542.
- [15]X.Li,Y.Shi and I.Gutman, *Graph Energy*, Springer, New york Heid.Dord., London 2012.
- [16]H.Liu,M.Lu and F. Tian, *Some upper bounds for the energy of graphs*, J.Math. Chem.,**411**(2007),45-57.
- [17]B.J.Mc Clelland,*Properties of the latent roots of a matrix: The estimation of  $\pi$ -electron energies*, J.of Chem.Phys., **54**(1971),640-643.
- [18]E. Sampathkumar, *The global domination number of a graph*, J. Math. Phys. Sci., **23**(1989),377-385.
- [19]E.Sampathkumar and L. Pushpa, *The global set-domination number of a graph*, Indian J.Pure appl.math. 2510,(1994),1053-1057.

- [20]S. K. Vaidya and R. M. Pandit, *Some new results on global dominating sets*, ISRN Dis. Math.,vol. 2012, ArticleID 852129, 6 pages, 2012. doi:10.5402/2012/852129.
- [21]S.K.Vaidya and R.M.Pandit, *Some results on global dominating sets*, Pro.J.Math. 323(2013),235-244.