

International Multidisciplinary Research Journal

Golden Research Thoughts

Chief Editor
Dr.Tukaram Narayan Shinde

Publisher
Mrs.Laxmi Ashok Yakkaldevi

Associate Editor
Dr.Rajani Dalvi

Honorary
Mr.Ashok Yakkaldevi

Golden Research Thoughts Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

Regional Editor

Manichander Thammishetty

Ph.d Research Scholar, Faculty of Education IASE, Osmania University, Hyderabad

International Advisory Board

Kamani Perera Regional Center For Strategic Studies, Sri Lanka	Mohammad Hailat Dept. of Mathematical Sciences, University of South Carolina Aiken	Hasan Bakfir English Language and Literature Department, Kayseri
Janaki Sinnasamy Librarian, University of Malaya	Abdullah Sabbagh Engineering Studies, Sydney	Ghayoor Abbas Chotana Dept of Chemistry, Lahore University of Management Sciences[PK]
Romona Mihaila Spiru Haret University, Romania	Ecaterina Patrascu Spiru Haret University, Bucharest	Anna Maria Constantinovici AL. I. Cuza University, Romania
Delia Serbescu Spiru Haret University, Bucharest, Romania	Loredana Bosca Spiru Haret University, Romania	Ilie Pinteau, Spiru Haret University, Romania
Anurag Misra DBS College, Kanpur	Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Xiaohua Yang PhD, USA
Titus PopPhD, Partium Christian University, Oradea,Romania	George - Calin SERITAN Faculty of Philosophy and Socio-Political Sciences Al. I. Cuza University, IasiMore

Editorial Board

Pratap Vyamktrao Naikwade ASP College Devrukh,Ratnagiri,MS India Ex - VC. Solapur University, Solapur	Iresh Swami S. D. M. Degree College, Honavar, Karnataka	Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur
R. R. Patil Head Geology Department Solapur University,Solapur	N.S. Dhaygude Ex. Prin. Dayanand College, Solapur	R. R. Yalikal Director Managment Institute, Solapur
Rama Bhosale Prin. and Jt. Director Higher Education, Panvel	Narendra Kadu Jt. Director Higher Education, Pune	Umesh Rajderkar Head Humanities & Social Science YCMOU,Nashik
Salve R. N. Department of Sociology, Shivaji University,Kolhapur	K. M. Bhandarkar Praful Patel College of Education, Gondia	S. R. Pandya Head Education Dept. Mumbai University, Mumbai
Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai	Sonal Singh Vikram University, Ujjain	Alka Darshan Shrivastava Shaskiya Snatkottar Mahavidyalaya, Dhar
Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College, Indapur, Pune	G. P. Patankar S. D. M. Degree College, Honavar, Karnataka	Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore
Awadhesh Kumar Shirotriya Secretary,Play India Play,Meerut(U.P.)	Maj. S. Bakhtiar Choudhary Director,Hyderabad AP India.	S.KANNAN Annamalai University,TN
	S.Parvathi Devi Ph.D.-University of Allahabad	Satish Kumar Kalhotra Maulana Azad National Urdu University
	Sonal Singh, Vikram University, Ujjain	

Golden Research Thoughts

GRT

ZOOPLANKTON DIVERSITY AND THEIR SEASONAL VARIATIONS OF KHANAPUR TANK, SHAHPUR, YADGIR DISTRICT, KARNATAKA

**Nagbhushan Reddy**

Environmental Biology Research Laboratory, Department of Zoology,
Sharnbasveshwar College of Science, Kalaburagi.




Co - Author Details :

Ramakrishna Reddy. B.¹, Siddaram .L.¹ and Vijayakumar. K²

¹Environmental Biology Research Laboratory, Department of Zoology,
Sharnbasveshwar College of Science, Kalaburagi.

²Department of Zoology, Gulbarga University, Kalaburagi , Karnataka.



ABSTRACT

The present work aims to study the seasonal variations and diversity of zooplankton in Khanapur Tank, Shahpur, Yadgir district for a period of one year from January 2014 to December 2014. Zooplankton community is also correlated with physicochemical parameters. Both qualitative and quantitative analysis of were conducted by taking samples from five sampling stations of the tank. We have recorded 30 species of which, 12 species belong to rotifera, 8 to cladocera, 6 to copepoda and 4 to ostracoda. Zooplanktons are heterotrophic planktonic animals floating in water. They serve as good indicators of changes in water quality. This study also reveals that different groups of zooplanktons have their own

peak periods of density, which is affected by local environmental conditions prevailing at that time.

KEYWORDS :Zooplanktons, Diversity, Seasonal variations, Khanapur Tank, Rotifera, Cladocera.

INTRODUCTION:

Zooplanktons are the smallest organisms present in almost all the water body and they can be observed only through microscope. They invariably form an integral component for fresh water communities and contribute significant to biological productivity. Zooplankton acts as main sources of food for many fishes and plays an important role in early detection and monitoring the pollution of

water. Zooplankton community distribution depends on some of the complex factors viz, change of climatic conditions, physical and chemical parameters and vegetation. Most of the planktonic organisms are cosmopolitan in distribution. A number of studies has been carried out on the condition of ecology and freshwater bodies in various parts of India but in some parts of southern Karnataka, the ecological studies of freshwater bodies especially zooplankton studies is very scanty. So that the present investigation made an attempt to study the zooplankton species in Bhadra Reservoir.

The study of freshwater fauna especially zooplankton even of a particular area is extensive and complicated due to environmental, physical, geographical and chemical variations involving ecological, extrinsic and intrinsic factors. This is particularly so with freshwater fauna (zooplankton), which plays a key role in preservation and maintenance of ecological balance. Its basic study is wanting and is absolutely necessary. Zooplankton exhibits a bimodal oscillation in a Spring and Autumn in the temperate lakes and reservoirs (Wetzel 2001). This fluctuation is greatly influenced by variations in the temperature along with many other factors. Among several factors, temperature seems to exhibit the greatest influence on the periodicity of zooplankton (Byars 1960; Battish and Kumari 1986; Prasad and Singh 2002). Water temperatures between 10–29°C are suitable for zooplankton development (Kaushik et al. 1992).

Zooplankton supports the economically important fish population. They are the major mode of energy transfer between phytoplankton and fish. A number of workers such as Ayyappan & Gupta (1980) Chakravarty (1985), Balkhi et al (1987) have reported on different aspects of zooplankton inhabiting Indian fresh waters.

The zooplankton community is composed of both primary consumers (which eat phytoplankton) and secondary consumers (which feed on the other zooplankton). They provide a direct link between primary producers and higher trophic levels like fish. Nearly all fish depend on zooplankton for food during their larval phases and some fishes continue to eat zooplankton throughout their entire lives (Madin et al., 2001).

MATERIAL AND METHODS

Study area:


Khanapur is one of the Village in Shahapur Taluk in Yadgir District in Karnataka State. It is a perennial fresh water body. It lies between 77.13 °E Longitude and 16.77 °N Latitude. Its water spread area is 110.3738 hectares. The Water of this water body too is used mainly for agriculture. In addition to this, its water is also used for washing clothes, bathing the animals and other domestic activities.



Location map of Yadgir District in the State



Map of Yadgir District Locating Khanapur



A satellite Map of Khanapur Tank



A view of General information of Khanapur Tank

Zooplankton sample Collection, Preservation, Identification and Density Analysis:

Water samples were collected from five stations in different locations of the tank during an early hours of the day (7 am to 9 am) to a period of 1 year (January 2014– December 2014) and such samples were pooled together to consider a final sample for analysis. The plankton net is made by the bolting nylon silk (mesh- size 50µm) is used for collection of zooplankton and which is conical shape and reducing cone with the bottle at its end. For a precise collection of zooplankton, the plankton net is towed horizontally and obliquely (for Qualitative) in surface water of the study area and for quantitative analysis, 10 liters of water samples were filtered out through the net. After transferring the sample in air tight plastic bottles, it would keep carefully with labeling and preserved immediately using 4% formaldehyde. Later, the collected samples were brought to laboratory and analyzed qualitatively under the microscope for different types of zooplanktons and identified using various authenticated

monographs 7- 11. After an accurate identification of each genus, the density of zooplankton was calculated as per the Lackey's drop method.

SEPARATION

In the laboratory the plankton collected from the tank were poured into a broad petridish. The different planktons were isolated with a pointed painting brush using a binocular dissection microscope. For qualitative and quantitative analysis of zooplankton, collections were made using a modified Haron-Trantor net with a square metallic frame of 0.0625 m² area. The filtering cone was made up of nylon bolting silk plankton net (No. 25 mesh size 50 μ). This net was used for collecting zooplankton. The net was hauled for a distance of 10 m. A collected sample was transferred to labeled vial containing 5% formalin. After sedimentation, 100 ml of sample was subjected to centrifugation at 1,500 rpm for 20 min and used for further investigation. In the laboratory the plankton were poured into a broad petri dish. The different planktons were isolated with a pointed painting brush using a binocular dissection microscope. Counting of the rotifera, cladocera, copepod and ostracoda was carried out using a Sedgewick Rafter cell. The data was used only to express the relative abundance of the principle zooplankton groups (Needham and Needham 1962).

RESULTS AND DISCUSSION

The obtained zooplankton forms were represented by four groups of Phylum viz Rotifera, Cladocera, Copepoda and Ostracoda. Among these, Rotifera comprise of 12 species, Cladocera 8 species, Copepoda 6 species, Ostracoda 4. All the dominant group of zooplankton were present throughout the year.

Rotifers: These are considered as the most important soft bodied invertebrates and they play a major role in aquatic food chain and major food for fishes. In rotifers zooplanktons like *Brachionous calcyflorus* and *Brachionous caudatus* found to be present throughout the year whereas cladocera was represented by *Bosmina longirostris*, *Alona davidi davidi*, *Macrothrix spinosa*, *Simocephalus exspinosus* and *Diaphanosoma species*. According to observation the Brachionus species are very common in temperate and tropical waters.

Cladocerans: Cladocera is an order of small crustaceans commonly they are called by "water fleas". It has been reported that the density and biomass of cladocerans was primarily determined by food supply. In the present study, similar observations were made where cladocerans were abundant when the food supply (phytoplankton) was maximum.

Copepods: These constitutes a major zooplankton communities occurring in almost all the water bodies, which serve food for many fish and play a vital role in ecological pyramids. The important factors which controlled the distribution of copepods were rainfall, river discharge and decreased phytoplankton abundance due to increased turbidity. In our studies the copepods recorded were *Cyclops strenuous*, *Cyclops scutifer* and *mesocyclops hyalinus*.

Ostracods: The ostracods are bivalve crustaceans found in both fresh and marine water. In the present study, only two species of ostracods were found and these are represented by Cypris species and heterocypris species.

The population of zooplankton in Khanapur tank consist of Rotifera, Copepoda, Cladocera, and

Ostracoda . The total number recorded were 2809 per liter of which rotifera 1,061 (37.44%), Cladocera 932 (33.17%), Copepoda 553 (19.69), and Ostracoda 263 (9.37%). All the above mentioned zooplanktons were dominant throughout the year. Diversity analysis shows that rotifers have 12 species, Cladocera 8, Copepods 6, and Ostracoda 4. The diversity of zooplankton groups is mentioned in Table 1.

High richness of rotifer species was also recorded during these seasons in contrast to the findings of Ayoagui & Bonecker (2004). This may be ascribed to the low population of cladocerans and the relief of the rotifers from competitive suppression by the cladocerans. Aoyagui & Bonecker (2004) stated that zooplankton diversity can be increased by the removal of competitively dominant species.

Table 1: Composition of zooplanktons

Months	Zooplankton groups			
	Rotifera	Cladocera	Copepoda	Ostracoda
January 2014	144	298	124	136
Feb	506	298	616	262
March	1732	1544	1056	182
April	1907	379	1468	588
May	1959	648	1193	706
June	1972	1478	943	432
July	1565	1728	200	196
August	1313	500	649	328
Sept	1270	1660	156	73
Oct	164	896		78
Nov	144	298	124	136
December 2014	137	902	130	84
Total	12726	11181	6637	3159
% of Plankton Diversity	37.44	33.17	19.69	9.37

CONCLUSION :

Among the zooplanktons in Khanapur tank, 30 species belong to Rotifera, Cladocera, Copepoda and ostracods. The overall view in this study reveals that the fluctuation of zooplankton occurs distinctly in the study area and normally in rainy season there is a less population due to the dilution factors and its effects leads to less photosynthetic activity by primary producers. The population raises a bit higher level during winter season due to favorable environmental conditions and presence of excess of food in the form of bacteria and suspended detritus, but in summer where inflow is less to compare with other seasons resulted in stability of water body and availability of food is more due to decomposition of organic matter and the density of zooplankton might be high due to less predators.

REFERENCES

- 1.Gannon, J.E., and Stemberger, R.S, Zooplankton especially crustaceans and rotifers as indicators of water quality. *Trans. Am. Micros. Soc.*, [97 (1978) 16-35].
- 2.Santhanam,P. and Perumal,P, Diversity of zooplankton in Parangipettai coastal waters,Southeast coast of India. *J.Mar.Biol.Assoc.India*, [45 (2003)144-151].

3. Padmavati, G. and Goswami, C, Zooplankton ecology in the Mandovi-Zuari estuarine system of Goa, west coast of India. *Int. J. Ecol. Environ. Sci.*, [31(1996) 273-278].
4. Arora, J. and Mehra, N, Seasonal dynamics of rotifers in relation to physical and chemical conditions of the river Yamuna (Delhi), India. *Hydrobiologia*, [491(2003) 101-109].
5. Rocha O., Matsumura-Tundisi, T., Espindola, E.L.G., Roche K.F., and Rietzler A.C. Ecological theory applied to reservoir zooplankton. In: *Theoretical reservoir ecology and its application* (Eds.: J.G. Tundisi and M. Straskraba). *Internat. Inst. Ecol.*, Sao Carlos, 457-476
6. Neves I. F., Rocha O. Roche K.F. and Pinto A.A., Zooplankton community structure of two marginal lakes of river Cuiaba (Mato Grosso, Brazil) with analysis of Rotifers and Cladocera diversity, *Braz. J. Biol.* 63 (33). Mukherjee B., *Environmental Biology*, Tata McGraw Hill Publishing Company Limited, New Delhi
7. Sinha B. and Islam M.R., Seasonal variation in zooplankton population of two lentic bodies and Assam State Zoo cum Botanical garden, Guwahati, Assam *Environ. Cons.*, 8, 273-278 (2002)
8. Singh, S.P., D. Pathak and R. Singh: Hydrobiological studies of two ponds of Satna (M.P), India *Cons.*, 8, 289-292 (2002)
9. Sharma, B.K. & Sharma, S. Freshwater Rotifers (Rotifera: Eurotatoria). In: *State Fauna Series: Zoological Survey of India*, Calcutta
10. Pathak S. K. and Mudgal L. K., A preliminary survey of zooplankton of Virla Reservoir of Khargaon. (Madhya Pradesh) India. *Indian, J. Env. &* (2002)
11. Arora, J. and Mehra, N. Seasonal dynamics of Rotifers to physical and chemical conditions of river Yamuna (Delhi), India, *Hydrobiologia*, 491, 101
12. Kiran B.R., puttaiah E.T and Kamath seasonal fluctuation of zooplankton in fish pond of Bhadra fish farm, Karnataka, *J. Zoos Print*
13. Schindler D.W. & Noven. B., Vertical distribution and seasonal abundance of zooplankton in two shallow lakes of the Experimental Lakes Area, northwestern Ontario *Fish. Res. Board Can.*, 28, 245
14. Saksena, D N. & Sharma S.P., Zooplankton fauna of some lentic water of Gwalior. Govinda sagar, Chhattri tank, Cladocera Copepoda Ostracoda Protozoa
15. Uttangi J.C.: Conservation and management strategy for the water fowls of minor irrigation tank habitats and their importance as stopover site in the Dharwad district, *Trends in wildlife and management*, Daya Publishing house, New Delhi, India: 179-221 (2001)
16. Hashemzadeh Farshad and G. V. Venkataramana: Impact of Physico-Chemical Parameters of Water on Zooplankton Diversity in Nanjangud Industrial Area, India, *Int. Res. J. Environment Sci.* Vol. 1(4), 37-42, (2012)
17. Thirupathaiah M., Sravanthy .Ch and sammaiah. Ch: Diversity of zooplankton in lower manair reservoir, Karimnagar, AP India, *I. Res. Biological Sci.*, 1(7), 27-32 (2012)
18. Sharma V., Bhoopendra Kumar V., and Madhu Sudan Sharma : Zooplankton Fauna in Relation to Physico-Chemical Characteristics in Madar Tank, Udaipur, Rajasthan, India,

Publish Research Article

International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication, you will be pleased to know that our journals are

Associated and Indexed, India

- * International Scientific Journal Consortium
- * OPEN J-GATE

Associated and Indexed, USA

- EBSCO
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Database
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Golden Research Thoughts
258/34 Raviwar Peth Solapur-413005, Maharashtra
Contact-9595359435
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com
Website : www.aygrt.isrj.org