

International Multidisciplinary
Research Journal

Golden Research
Thoughts

Chief Editor
Dr.Tukaram Narayan Shinde

Publisher
Mrs.Laxmi Ashok Yakkaldevi

Associate Editor
Dr.Rajani Dalvi

Honorary
Mr.Ashok Yakkaldevi

Welcome to GRT

RNI MAHMUL/2011/38595

ISSN No.2231-5063

Golden Research Thoughts Journal is a multidisciplinary research journal, published monthly in English, Hindi & Marathi Language. All research papers submitted to the journal will be double - blind peer reviewed referred by members of the editorial board. Readers will include investigator in universities, research institutes government and industry with research interest in the general subjects.

International Advisory Board

Flávio de São Pedro Filho Federal University of Rondonia, Brazil	Mohammad Hailat Dept. of Mathematical Sciences, University of South Carolina Aiken	Hasan Baktir English Language and Literature Department, Kayseri
Kamani Perera Regional Center For Strategic Studies, Sri Lanka	Abdullah Sabbagh Engineering Studies, Sydney	Ghayoor Abbas Chotana Dept of Chemistry, Lahore University of Management Sciences[PK]
Janaki Sinnasamy Librarian, University of Malaya	Ecaterina Patrascu Spiru Haret University, Bucharest	Anna Maria Constantinovici AL. I. Cuza University, Romania
Romona Mihaila Spiru Haret University, Romania	Loredana Bosca Spiru Haret University, Romania	Ilie Pinteau, Spiru Haret University, Romania
Delia Serbescu Spiru Haret University, Bucharest, Romania	Fabricio Moraes de Almeida Federal University of Rondonia, Brazil	Xiaohua Yang PhD, USA
Anurag Misra DBS College, Kanpur	George - Calin SERITAN Faculty of Philosophy and Socio-Political Sciences Al. I. Cuza University, IasiMore
Titus PopPhD, Partium Christian University, Oradea, Romania		

Editorial Board

Pratap Vyamktrao Naikwade ASP College Devrukh, Ratnagiri, MS India	Iresh Swami Ex - VC. Solapur University, Solapur	Rajendra Shendge Director, B.C.U.D. Solapur University, Solapur
R. R. Patil Head Geology Department Solapur University, Solapur	N.S. Dhaygude Ex. Prin. Dayanand College, Solapur	R. R. Yalikal Director Management Institute, Solapur
Rama Bhosale Prin. and Jt. Director Higher Education, Panvel	Narendra Kadu Jt. Director Higher Education, Pune	Umesh Rajderkar Head Humanities & Social Science YCMOU, Nashik
Salve R. N. Department of Sociology, Shivaji University, Kolhapur	K. M. Bhandarkar Praful Patel College of Education, Gondia	S. R. Pandya Head Education Dept. Mumbai University, Mumbai
Govind P. Shinde Bharati Vidyapeeth School of Distance Education Center, Navi Mumbai	G. P. Patankar S. D. M. Degree College, Honavar, Karnataka	Alka Darshan Shrivastava Shaskiya Snatkottar Mahavidyalaya, Dhar
Chakane Sanjay Dnyaneshwar Arts, Science & Commerce College, Indapur, Pune	Maj. S. Bakhtiar Choudhary Director, Hyderabad AP India.	Rahul Shriram Sudke Devi Ahilya Vishwavidyalaya, Indore
Awadhesh Kumar Shirotriya Secretary, Play India Play, Meerut (U.P.)	S. Parvathi Devi Ph.D.-University of Allahabad	S. KANNAN Annamalai University, TN
	Sonal Singh, Vikram University, Ujjain	Satish Kumar Kalhotra Maulana Azad National Urdu University

Address:- Ashok Yakkaldevi 258/34, Raviwar Peth, Solapur - 413 005 Maharashtra, India
Cell : 9595 359 435, Ph No: 02172372010 Email: ayisrj@yahoo.in Website: www.aygrt.isrj.org

COMPARATIVE STUDY OF BEHAVIORAL RESPONSE DUE TO ACUTE EXPOSURE OF MALATHION AND CYPERMETHRIN ON FRESHWATER FISH, *LEPIDOCEPHLICTHYS GUNTEA* (HAMBUCH).



Patil M. U.

Dept. Of Zoology, V.U.Patil Arts, and Late Dr. B. S. Desale science college, Sakri, Dist-Dhule

Short Profile

Patil M. U. is a Head at Department of Zoology in V. U. Patil Arts, and Late Dr. B. S. Desale science college, Sakri, Dist-Dhule. She has completed B.Ed., M.Sc., M. Phil., M.Z.S.I. She has professional experience of 7 years and research experience of 9 years. She has done one minor research project.

Co-Author Details :

S. S. Patole

Dept of Zoology, V,V,M's S.G.Patil Arts Science and Commerce College, Sakri, District-Dhule .

K. D. Ahirrao

Head, Dept of Zoology, Rani Laxmibai College, Parola, Dist-Jalgaon .



ABSTRACT:

Aquatic contamination by pesticides is arising as a result of their extensive use in agriculture and public health programmes. However, indiscriminate use of these pesticides for crop protection causes much damage to the non-target animals. The aquatic organisms particularly fish becomes highly sensitive, as water gets contaminated by toxic chemicals, some pesticides are not easily degraded in water and fishes are

known to be absorbing them through the skin in high quantities. They affect the growth, metabolism and on physical and physiological activities and ultimately on the survival of the fish. In the present study pesticides, Malathion and Cypermethrin are highly toxic to fish *Lepidocephalichthys guntea* at acute exposure leading the effect on several behaviors. But the synergistic effects are more harmful than any above pesticide.

KEYWORDS

L. guntea, behavior, Malathion, Cypermethrin,

1. INTRODUCTION:

Pesticides have been recognized as serious pollutants of the aquatic ecosystems with drastic effect to aquatic fauna (Sambasiva, 1999). Several workers worked on the fresh water fishes diversity (Jaiswal and Ahirrao, 2012; Ahirrao, 2014) but fish behavior and impact of pesticides and environmental pollutants are not understood properly in an aquatic organism. Chemical pesticides when applied on the field affect the survival, growth, metabolism and on physiological activities on non-target organisms (Murugesen and Haniffa, 1992). The aquatic organisms especially fish becomes highly sensitive in such ecosystem. When the water gets contaminated by toxic chemicals, some pesticides are not easily degraded in water and fishes are known to be absorbing them through the skin in high quantities (Tilak *et. al.*, 2001). Toxicants may affects the aquatic organisms in many ways and categorized their effects as direct and indirect (Bhaskar *et. al.*, 2002). These pesticides are not readily removed by usual excretory routes, because they inhibits by the enzymes participating in the detoxification and excretory reaction of fishes (Kulshrestha *et. al.*, 1984). Pesticides are known to have hazardous effect on various organ systems of fish. They also alter the cellular morphology and physiology of fishes. The fishes get sicker and killed when pollution develop in the water bodies. However, in that situation they may deliberately leave the polluted area, but sudden pollution of rivers and streams result in the death of fishes (Sahai, 1988).

MATERIALS AND METHODS

The fresh water Fish, *Lepidocephlichthys guntea* were collected from 'Panzara Kan River' around Sakri, district Dhule, Maharashtra State, India. They were brought to the laboratory, cleaned and were placed in well aerated glass aquaria containing sufficient water for fifteen days for acclimatization. They were fed with fish diet. The healthy, active, mature adults, approximately same sized (4.0 + 0.5 cm length) and weighing about 10.0 + 0.3 grams were exposed to pesticide to calculate LC₅₀ values for pesticides viz. Organophosphate (Malathion) and Synthetic Pyrethroid (Cypermethrin). Values are estimated by the method of Probit Analysis (Finney, 1971). The corrected present mortality was calculated by applying Abbott's formula (1945).

Static-with-renewal acute toxicity tests were conducted with ten fish in each graded concentration. Fish were placed in five glass aquaria containing dechlorinated tap water. The physic chemical parameters of water were carried out according to APHA, (1998). Before to starting of experimentation and values are as follows:

PH = 7.1 + 1.0 ; Temperature: 26.0 + 2.0°C ; Dissolved Oxygen : 5.0 to 5.5 mg/l.; Free CO₂ = 2.9 to 3.1 mg/l.; Alkalynity(CO₃)= 33 to 35 mg/l ; Alkalynity(HNO₃)=380 to 390 mg/l ; Chloride=540 to 550 mg/l ; Sulphate 10.50 to 11.0 ppm ; Hardness=145 to 160 mg/l

Thereafter, Malathion and Cypermethrin were added as per the ascending series of various concentrations (1 to 20 ppm). The fish kept in chlorine free tap water served as a control. Food was withheld 24 hours prior to acute toxicity tests. The test solution was replaced and mortality monitored at 24, 48, 72 and 96 hours. Fishes were considered dead if they don't respond to stimulus touching with glass rod. Dead fish were removed immediately and the observations were recorded. The entire experiment was repeated five times. Behavioral changes of the experimental fishes are recorded during the exposure period upto 96 hours. Fishes were exposed to predetermined LC₅₀ Values of Malathion (11 ppm) and Cypermethrin (5.2 ppm) for 96 hours.

OBSERVATIONS AND RESULTS:

Table:1. Behavioral changes during intoxication in experimental fish, *L. guntea* observed with exposure to Malathion, Cypermethrin and Synergistic effects of Malathion and Cypermethrin.

Behavior	Observations	Control	Malathion	Cypermethrin	Synergistic effects	
					3/4 th Mal.+ 1/4 th Cyp	1/4 th Mal +3/4 th Cyp
Mucous Secretion	More mucous secretion takes place	Normal secretion of the mucous all over the body of <i>L. guntea</i>	+	++	+	+++
Respiratory activity	Repeatedly opening and closing of mouth because of cough	Normal opening and closing of mouth	+	++	++	+++
Yawn	Wide opening of mouth	Normal movement of jaws of mouth	++	++	++	++
Body Color	Changes occur due to pesticide	It is bright brownish at dorsal side and milky white at ventral side	+	++	++	+++
Swimming Activities	Frequently come to the water surface and erratic swimming	Swimming normally	+	++	++	+++
Jerking movements	S jerk: Movement of body sequentially from head to tail	No jerking movements	+	+	++	+++
	Partial jerk: Movement of Head or tail	No jerking movements	+	+++	+++	+++
	Bending body	Normal movements	+	+	+++	++
Abnormal behavior	Dash to the wall of an aquaria	Not dashed even though they may be overcrowded	+	++	+	+++
	Jumping out from aquaria	Generally this behavior is not found.	+	++	++	+++
Aggressive behavior	Nudge (Movement of fish towards another fish)	Normally move	++	++	++	+++
	Nip (Biting to another fish)	Don't nip	++	++	++	+++
Gill size	Swelling at gill side and it gets reddish in color	Normal shape size and color of gill	+	++	++	+++
Equilibrium	Lost Equilibrium	Do not loss equilibrium and all movements are found normally	+	++	+	+++
	Paralysis	Never suffering from paralysis	++	++	++	+++
	Death	Survive normally	+	+	++	+++

Values from table: + = less number of observations recorded (Hypo activity)
 ++ = Moderately recorded observations
 +++ = More activities recorded (Hyperactivity).

DISCUSSION:

During pesticide treatment, fishes responded by expressing changes of their physical mode of behavior. These behavioral responses of fishes varied in accordance with predetermined sub-lethal concentration of pesticides viz. Malathion, Cypermethrin and synergistic effects of Malathion and Cypermethrin at concentration of 3/4th Mal. + 1/4th Cyp and 1/4th Mal + 3/4th Cyp at 96 hours to experimental fish *L. guntea*.

Lethal concentrations are gaining importance now a day. This is so because distinct behavioral changes are infested under sub-lethal exposure considering the amount of pesticide used in field application (Patil and Patole, 2011; Ganesan, et al., 1989;). It is lost degradation, spill over, runoff etc. It is ultimate mixing to the rivers, tanks and irrigation canals and of the volume of water present in these stronger systems. It is more appropriate that inhabitants of the fresh water ecosystem are more likely to expand to the low concentration of pesticides than to high concentration. Beside pointed that most of

the pesticides are relatively toxic even at low concentrations, from all these studies under sub-lethal concentration. Hence it is selected for the present study.

Similar observations have been reported by earlier researchers in fishes intoxicated by different pesticides in different species at different period of exposure. The behavioral changes in the manifestation of the motivational, biochemical (Patil and Patole 2012), physiological and environmentally influenced state of organism. They decreased opercular movement and corresponding increased in frequency of surfacing of fish clearly indicates that fish adoptively shift towards aerial respiration(by obtaining atmospheric oxygen surfacing) and the fish tries to avoid contact with the pesticides through gill chamber (Santkumar and Balaji, 2000). The erratic swimming of the treated fish indicated loss of equilibrium. It is likely that the region in the brain. Which is associated with the maintenance of the equilibrium should have been affected (Rao and Rao, 1987). The erratic swimming jerky movement and the convulsions before death were evident and the severity varied with pesticide concentration indicates the sign of asphyxiation as indicated by death (Kalavathy *et. al.*, 2001). The surfacing phenomenon of fish observed under cypermethrin exposure might be due to hypoxic condition of the fish as reported by Radhaiah and Jayntha Rao (1988). The increased surfacing during the initial period of exposure to Cypermethrin and Malathion concentrations suggest and elevated rate of metabolism changes in ventilation rate and changes in surfacing frequencies are the general symptoms noticed in the fish after exposure to pesticide and these activities help the fish to avoid contact with poison and fight against stress (Jain, 2001).

Aggressive behavior such as nudge and nip were increased following exposure to the toxic material. Orientation and locomotory pattern were found to be involved in most aspects of fish behavior such as migration, feeding, mating and courtship. Which were altered under stress conditions of environmental toxicants (Shanthakumar and Balaji, 2000; Prasad *et al.*, 2002). The accumulation of increased secretion of mucous in the fish exposed to Cypermethrin may e adoptive response perhaps providing additional protection against corrosive nature of the pesticide and to avoid the absorption of the toxicant by the general body surface. Jagdish and Sahai (1989) reported at higher concentration of pesticides, *Heteropneustes fossilis* and *Mystus vittatus* exhibited rapid opercular movement as well as erratic swimming. They often lost their balanced dashed against the wall of aquaria and even tried to escape out. Jyoti *et al.*,(1989) observed increased opercular movements in *Channa punctatus* on administration of pesticides , Malathion. The body color of the treated fish become pale on prolonged exposure the movements of the fish were greatly arrested and opercular movement shows down. Khillare(1990) observed negative response to the stress environment in *Channa gachua* exposed to Dimethoate and Rogor. Normal activities of the fin movements and the response o the stimuli were very much affected. We agree with the earlier findings because same observations were found in present studies.

CONCLUSION:

In the present study it has been found that both pesticides i.e. Malathion and Cypermethrin are highly toxic to fish *Lepidocephalichthys guntea* at acute exposure leading the effect on several behaviors of fish as well as entire physiological studies are found to be sensitive change in the physiological indicators which reflects change in normal activities of various functional systems. The synergistic effects are more harmful than that of the individual higher concentration of pesticide. The random use of pesticide must be avoided for pressuring aquatic biota.


REFERENCES:

1. Abbott, W.S. (1925). Methods of computing the effectiveness of an insecticide, *J. Econ. Ent.* 8:265-267.
2. Ahirrao, K. D. (2014). Fish diversity of the Bori dam at Tamaswadi, Parola, district Jalgaon, Maharashtra state. *Golden research thoughts, Vol.3(12)*1-5.
3. APHA/AWWA/WEF (1998). Standard method for the evolution of water and waste water. 20th edition, *American Public Health Association*, New York, USA.
4. Bhaskar, P., Murthy, E. and Narain, E. (2002). Toxicity of Diamethoate and Monocil on *Channa marulius*. *J. Environ. Biol.*, 20(5) : 319-324.
5. Finney, D.J. (1971). Probit analysis. *Cambridge University Press*, London.
6. Ganesan, R.M., Jebakumar, S. R. D., and Jayraman, J.(1989). Sublethal effect of Organochlorin insecticides Endosulfan on Protein, Carbohydrate and Lipid content on liver tissue of *Oreochromis mossambicus*, *Proc. Indian Acad. Sci. (Anim. Sci.)*98 (1): 51-52.
7. Jain, G.K. (2000). Pesticides and fish, *Aditya Publishurs*, Bina (M.P.)
8. Jaiswal, D. P. and K. D. Ahirrao (2012). Ichthyodiversity of the Rangawali dam, Navapur, district Nandurbar, Maharashtra state. *Journal of of research in biology. Vol-2(3)*:241-245.
9. Jyoti, A., Khan, A. A., Haque N., Masroor, F., Barbhauyah, S. I. (1989). Studies on the effect of Malathion on a freshwater fish, *Channa punctatus*. *J. Envio., Biol.* 10(3): 251-257.
10. Kalavathy, K. A., Sivakumar, A. and Chandran, R. (2001). Toxicity of pesticide Diamethoate on the fish, *Sarotherodon mossambicus*. *J. Eco. Res. Biocn.*, 2(2):27-32.
11. Khillare, Y. K. (1990). Effect of pesticide on freshwater fish, *Puntius stigma*. In *Trend in Ecotoxicol. Acad. Environ. Biol.* Karad, 185-188.
12. Kulshrestha, S. K., Srivastava R. S. and Arora, N. (1984). Toxicity of two pesticides to the Kidney of a freshwater *Channa striatus*. (Bloch). *Proc. Sem. Eff. Pest. Aq. Fau.*, 63-70.
13. Murugesan, A. G. and Haniff, M. A. (1992). Histopathological and Histochemical change in the Oocyte of the air breathing fish, *Heteropneustes fossilis* (Bloch) Exposed to textile mill effluent. *Bull. Environ. Contam. Taxicol.* 48: 929-936.
14. Patil, M.U. and Patole S.S., (2011). Studies on Effect of Insecticides on Oxygen Consumption of Fresh Water Fish, *Lepidocephalichthys guntea*(Ham-Buch) *Ecology and Fisheries*, Vol.4(2):29-32.
15. Patil M.U. and Patole S.S. (2012). Effect of Malathion and Cypermethrin on Biochemical Constituents of Freshwater Fish, *Lepidocephalichthys guntea*(Ham-buch) *International Indexed & Research Journal*, Vol IV,(39),33-35.
16. Prasad M., Bandopadhaya and Aditya, A. (2002). Xenobiotic impact on sensitivity in *Anabas testiduneus* (Bloch). *J. Eco. Biol.* 14: 11-124.
17. Radhaih, V. and Jayantha, Rao (1988). Behavioral changes in freshwater fish, *Tilipia mossumbicus* expose to Fenvalerate. *Environ. And Ecol.* 6:22-23.
18. Rao, B. and Rao. C. (1987). Independent and combine action of Carbaryl and Phenthoate on snake head fish, *Channa punctatus* (Bloch). *Curr. Sci.* 56(7):331-332.
19. Sahai, S. (1988). Pesticide accumulation in some tissues of *Rasbora doniconius* (Teleostei). *I:symp. On environmental impact on animal and aquaculture*, 33:657-659.
20. Sambashiva Rao, K. R. S. (1999). Pesticide impact on fish metabolism, *Discovery Publication House, New, Delhi*.
21. Santhakumar, M. and Balaji, M. (2000). Acute toxicity of an organophosphorous insecticide

Monochrotophos and its effect on its behavior of an air breathing fish, *Anabas testudineus*(Bloch). *J. Environ. Biol.* 21(2): 121-123.

22.Tilik, K. S., Veeriah, K. and Yacobu, K.(2001). Studies on Histopathological changes in the Gill, Liver and Kidney of *Ctenopharyngodon idella* (Valencinnes) exposed to technical Fenvalerate an EC_{20} . *Pollu. Res.* 20(3):387-393.

Map: Showing Collection site at Panzara Kan River near Sakri, Tal-Sakri, district-Dhule(MS), India.



Source: Map from www.google.com(India)

Publish Research Article

International Level Multidisciplinary Research Journal For All Subjects

Dear Sir/Mam,

We invite unpublished Research Paper, Summary of Research Project, Theses, Books and Book Review for publication, you will be pleased to know that our journals are

Associated and Indexed, India

- ★ International Scientific Journal Consortium
- ★ OPEN J-GATE

Associated and Indexed, USA

- EBSCO
- Index Copernicus
- Publication Index
- Academic Journal Database
- Contemporary Research Index
- Academic Paper Database
- Digital Journals Database
- Current Index to Scholarly Journals
- Elite Scientific Journal Archive
- Directory Of Academic Resources
- Scholar Journal Index
- Recent Science Index
- Scientific Resources Database
- Directory Of Research Journal Indexing

Golden Research Thoughts
258/34 Raviwar Peth Solapur-413005, Maharashtra
Contact-9595359435
E-Mail-ayisrj@yahoo.in/ayisrj2011@gmail.com
Website : www.aygrt.isrj.org